

Convergence of random variables, and the Borel-Cantelli lemmas

Lecturer: James W. Pitman

Scribes: Jin Kim (jin@eecs)

1 Convergence of random variables

Recall that, given a sequence of random variables X_n , almost sure (a.s.) convergence, convergence in P , and convergence in L^p space are true concepts in a sense that $X_n \rightarrow X$. In this lecture, we will define weak convergence, or convergence in distribution, $P_{X_n} \rightarrow P_X$, which we write, by abuse of notation, $X_n \xrightarrow{d} X$.


Definition 1.1 (Convergence in distribution) We say $X_n \xrightarrow{d} X$ if $P(X_n \leq x) \rightarrow P(X \leq x)$ for all x at which the RHS is continuous.

This weak convergence appears in the central limit theorem.

Theorem 1.2 $X_n \xrightarrow{d} X \iff E f(X_n) \rightarrow E f(X)$ for all bounded and continuous function f .

Proof See Durrett. \square

Theorem 1.3 The following property holds among the types of convergence.



Proof (*) can be proven by Chebychev inequality (with usually $p = 2$):

$$P(|X_n - X| > \epsilon) \leq \frac{E |X_n - X|^p}{\epsilon^p},$$

and $(**)$ is proven in Durrett. \square

Exercise. counter examples

- \xrightarrow{p} but not $\xrightarrow{\text{a.s.}}$: moving blip
- \xrightarrow{p} but not $\xrightarrow{L^p}$: try $X_n = n\mathbf{1}(0, 1/n)$. $X_n \xrightarrow{p} 0$ but $E|X_n - 0| = 1$, thus $X \not\xrightarrow{L^1} 0$ in L^1 .

Proposition 1.4 (Inducing a Metric) $X_n \xrightarrow{\text{a.s.}} X$ cannot be metrized, but $X_n \xrightarrow{L^p} X$ and $X_n \xrightarrow{p} X$ can be metrized, e.g. using $E(|X_n - X| \wedge 1)$. Furthermore, when so metrized, the space of random variables are complete.

Proof See text (uses BCL).

Definition 1.5 (Infinitely Often (i.o.) and Eventually (ev.)) Let q_n be some statement, e.g., $|X_n - X| > \epsilon$. We say $(q_n \text{ i.o.})$ if for all n , $\exists m \geq n$: q_m is true, and $(q_n \text{ ev.})$ if $\exists n$: for all $m \geq n$: q_m is true.

Exercise. Note that the following holds;

- $X_n \longrightarrow X \iff \forall \epsilon > 0, |X_n - X| < \epsilon \text{ ev.}$
- $X_n \not\longrightarrow X \iff \forall \epsilon > 0, |X_n - X| > \epsilon \text{ i.o.}$
- $(q_n \text{ i.o.})^\sim = (q_n \text{ ev.})$

Similarly, for a sequence of events A_n in a prob space (Ω, \mathcal{F}, P) , we can say the following;

- $(A_n \text{ i.o.}) = \{\omega : \omega \in A_n \text{ i.o.}\} = \bigcap_n \bigcup_{m \geq n} A_m$
- $(A_n \text{ ev.}) = \{\omega : \omega \in A_n \text{ ev.}\} = \bigcup_n \bigcap_{m \geq n} A_m$
- $(A_n \text{ i.o.})^c = A_n^c \text{ ev.}$

Main application of the idea of i.o. and ev. is to the proof of a.s. convergence. For example, since

$$(X_n \longrightarrow X) = \bigcap_{\epsilon > 0} (|X_n - X| < \epsilon \text{ ev.}) ,$$

we have

$$P(X_n \longrightarrow X) = \lim_{\epsilon \searrow 0} P(|X_n - X| < \epsilon \text{ ev.}) .$$

Since the basic criterion for a.s. convergence can be written as

$$(X_n \longrightarrow X) \iff \forall \epsilon > 0, P(|X_n - X| > \epsilon \text{ i.o.}) = 0 ,$$

we are interested in conditions in some sequence of events A_n so that $P(A_n \text{ i.o.}) = 0$.

2 Borel-Cantelli Lemma

Theorem 2.1 (Borel-Cantelli Lemma)

1. If $\sum_n P(A_n) < \infty$, then $P(A_n \text{ i.o.}) = 0$.
2. If $\sum_n P(A_n) = \infty$ and A_n are independent, then $P(A_n \text{ i.o.}) = 1$.

There are many possible substitutes for independence in BCL II, including Kochen-Stone Lemma.

Before proving BCL, notice that

- $\mathbf{1}(A_n \text{ i.o.}) = \lim_{n \rightarrow \infty} \sup \mathbf{1}(A_n)$
- $\mathbf{1}(A_n \text{ ev.}) = \lim_{n \rightarrow \infty} \inf \mathbf{1}(A_n)$
- $(A_n \text{ i.o.}) = \lim_{m \rightarrow \infty} P(\cup_{n \geq m} A_n)$ (as $m \uparrow$, $\cup_{n \geq m} A_n \downarrow$)
- $(A_n \text{ i.o.}) = \lim_{m \rightarrow \infty} P(\cap_{n \geq m} A_n)$ (as $m \uparrow$, $\cap_{n \geq m} A_n \downarrow$).

Therefore,

$$\begin{aligned}
 P(A_n \text{ ev.}) &\leq \lim_{n \rightarrow \infty} \inf P(A_n) && \text{by Fatou's lemma} \\
 &\leq \lim_{n \rightarrow \infty} \sup P(A_n) && \text{obvious from definition} \\
 &\leq P(A_n \text{ i.o.}) && \text{dual of Fatou's lemma (i.e. apply to } (\cdots)^\sim)
 \end{aligned}$$

Pf of BCL I

$$\begin{aligned}
 P(A_n \text{ i.o.}) &= \lim_{m \rightarrow \infty} P(\cup_{n \geq m} A_n) \\
 &\leq \lim_{m \rightarrow \infty} \sum_{n \geq m} P(A_n) = 0 \quad \text{since } \sum_{i=1}^{\infty} P(A_n) < \infty. \quad \square
 \end{aligned}$$

Pf of BCL I (Alternative method)

Consider a random variable $N := \sum \mathbf{1}(A_n)$, i.e. the number of events that occur. Then $E[N] = \sum_{n=1}^{\infty} P(A_n)$ by the Monotone Convergence Theorem, and

$$\begin{aligned}
 \sum_{n=1}^{\infty} P(A_n) < \infty &\implies E[N] < \infty \\
 &\implies P(N < \infty) = 1 \\
 &\implies P(N = \infty) = 0 \\
 &\implies P(A_n \text{ i.o.}) = 0 \quad \text{because } (N = \infty) \equiv (A_n \text{ i.o.}). \quad \square
 \end{aligned}$$

Pf of BCL II We will show that $P(A_n^c \text{ ev.}) = 0$.

$$P(A_n^c \text{ ev.}) = \lim_{n \rightarrow \infty} P(\cap_{m \geq n} A_m^c) = \lim_{n \rightarrow \infty} \prod_{m \geq n} P(A_m^c) \quad (1)$$

$$\begin{aligned} &= \lim_{n \rightarrow \infty} \prod_{m \geq n} (1 - P(A_m)) \leq \lim_{n \rightarrow \infty} \prod_{m \geq n} \exp(-P(A_m^c)) \\ &= \lim_{n \rightarrow \infty} \exp\left(-\sum_{m \geq n} P(A_m^c)\right) = 0 \end{aligned} \quad (2)$$

For (1) we used the following fact (due to the independence of A_n);

$$P(\cap_{m \geq n} A_m^c) = \lim_{N \rightarrow \infty} P(\cap_{n \leq m \leq N} A_m^c) = \lim_{N \rightarrow \infty} \prod_{n \leq m \leq N} P(A_m^c) = \prod_{n \leq m} P(A_m^c)$$

and $1 - x \leq \exp(-x)$ was used in (2). \square

As a trivial example, consider $A_n = (0, 1/n)$ in $(0, 1)$. Then, $P(A_n) = 1/n$, $\sum P(A_n) = \infty$, but $P(A_n \text{ i.o.}) = P(\emptyset) = 0$.

Intuitive example Consider random walk in \mathbf{Z}^d , $d = 0, 1, \dots$. $S_n = X_1 + \dots + X_n$, $n = 0, 1, \dots$ where X_i are independent in \mathbf{Z}^d . In the simplest case, each X_i has uniform distribution on 2^d possible strings. i.e., if $d = 3$, we have $2^3 = 8$ neighbors

$$\left\{ \begin{array}{c} (+1, +1, +1) \\ \vdots \\ (-1, -1, -1) \end{array} \right\}.$$

Note that each coordinate of S_n does a simple coin-tossing walk independently. We can prove that

$$P(S_n = 0 \text{ i.o.}) = \begin{cases} 1 & \text{if } d = 1 \text{ or } 2 \quad (\text{recurrent}) \\ 0 & \text{if } d \geq 3 \quad (\text{transient}). \end{cases} \quad (3)$$

Sketch of Pf of (3)

Let us start with $d = 1$, then

$$P(S_{2n} = 0) = \binom{2n}{n} 2^{-2n} \sim \frac{c}{\sqrt{n}} \text{ as } n \rightarrow \infty.$$

where we used the fact, $n! \sim \left(\frac{n}{e}\right)^n \sqrt{2\pi n}$.

Note

$$\sum \left(\frac{1}{\sqrt{n}}\right)^d = \begin{cases} \infty & d = 1, 2 \\ < \infty & d = 3, 4, \dots \end{cases} \quad (4)$$

BC II and (4) together gives (3). \square

Because

$$X_n \rightarrow X \text{ a.s.} \iff X_n - X \rightarrow 0 \text{ a.s.},$$

thus it is enough to understand as convergence to 0.

Proposition 2.2 *The following are equivalent:*

1. $X_n \longrightarrow 0$
2. $\forall \epsilon > 0, P(|X_n| > \epsilon \text{ i.o.}) = 0$
3. $M_n \longrightarrow 0$ where $M_n := \sup_{k \geq n} |X_k|$
4. $\exists \epsilon_n \downarrow 0 : P(|X_n| > \epsilon_n \text{ i.o.}) = 0$

If we need to show $X_n \xrightarrow{\text{a.s.}} X$ but do not know X , then it might be easier to show instead that $P(X_n \text{ is a Cauchy sequence}) = 1$. This leads to the following;

Lemma 2.3 *Let X_n be any sequence of random variables, and define $M_n := \sup_{m \geq n} |X_n - X_m|$. Then*

$$\exists X : X_n \longrightarrow X \text{ a.s.} \iff M_n \xrightarrow{p} 0$$

Proof Consider $M_n^* := \sup_{m, p \geq n} |X_m - X_p|$. Notice $M_n^* \downarrow$. Thus

$$M_n^* \xrightarrow{p} 0 \iff M_n^* \xrightarrow{\text{a.s.}} 0$$

Combine with the previous result to finish the proof. \square