

Probability

Introduction

There are two central questions concerning probability. First, what are its formal features? That is a *mathematical* question, to which there is a standard, widely (though not universally) agreed upon answer. This answer is reviewed in the next section. Second, what sorts of things *are* probabilities—what, that is, is the *subject matter* of probability theory? This is a *philosophical* question, and while the mathematical theory of probability certainly bears on it, the answer must come from elsewhere. To see why, observe that there are many things in the world that have the *mathematical structure* of probabilities—the set of measurable regions on the surface of a table, for example—but that one would never mistake for *being* probabilities. So probability is distinguished by more than just its formal characteristics. The bulk of this essay will be taken up with the central question of what this “more” might be.

Kolmogorov’s axiomatization

Probability theory was inspired by games of chance in seventeenth century France and inaugurated by the Fermat-Pascal correspondence, which culminated in the *Port-Royal Logic* (Arnauld, 1662). Its axiomatization had to wait nearly another three centuries. The *locus classicus* of the mathematical theory of probability is Kolmogorov (1933), who found his inspiration in measure theory. His axiomatization has become orthodoxy. Let Ω be a non-empty set. A *field (algebra)* on Ω is a set \mathcal{F} of subsets of Ω that has Ω as a member, and that is closed

under complementation (with respect to Ω) and union. Assume for now that \mathcal{F} is finite. Let P be a function from \mathcal{F} to the real numbers obeying:

1. $P(a) \geq 0$ for all $a \in \mathcal{F}$.
2. $P(\Omega) = 1$.
3. $P(a \cup b) = P(a) + P(b)$ for all $a, b \in \mathcal{F}$ such that $a \cap b = \emptyset$.

Call P a *probability function*, and (Ω, \mathcal{F}, P) a *probability space*.

One could instead attach probabilities to members of a collection of *sentences* of a formal language, closed under truth-functional combinations. Either way, a kind of reflective equilibrium is achieved between these axioms, which are thought to be intuitively plausible, and various important interpretations of probability (to be discussed in the subsequent sections), which obey them, and which bring them to life in applications.

It is often thought that the only non-conventional part of the axiomatization is the third axiom. That is too quick. For it is substantive that probabilities are:

- (i) defined by *functions* (rather than by one-many or many-many mappings);
- (ii) functions of *one variable* (unlike primitive conditional probability functions, which are functions of two variables);
- (iii) defined on a *field* (rather than a set with weaker closure conditions);
- (iv) represented *numerically* (rather than qualitatively, as ‘possibility’ is, or comparatively, as ‘similarity to a given world’ is in the Stalnaker/Lewis-style semantics for counterfactuals (Lewis [1986]));
- (v) *real numbers* (rather than those of some other number system);
- (vi) *bounded* (unlike other quantities that are treated measure-theoretically, such as lengths);
- (vii) attain *maximal* and *minimal* values (thus prohibiting open or half-open ranges).

For a discussion of rival theories that relax or replace (ii), (iii), (iv) and (vi), see Fine (1973). Complex-valued probabilities are proposed by Feynman and Cox (see Mückenheim et al. 1986); infinitesimal probabilities (of non-standard analysis) by Skyrms (1980) and Lewis (1980) among others; unbounded probabilities by Renyi (1970). Primitive conditional probability functions will be briefly discussed at the end of this section.

Kolmogorov extends his axiomatization to cover infinite probability spaces. Probabilities are now defined on a σ -field (σ -algebra)—a field that is further closed under *countable* unions—and the third axiom is correspondingly strengthened:

3'. (Countable additivity) If a_1, a_2, a_3, \dots is a countable sequence of (pairwise) disjoint sets, each belonging to \mathcal{F} , then

$$P\left(\bigcup_{n=1}^{\infty} a_n\right) = \sum_{n=1}^{\infty} P(a_n).$$

De Finetti (1990) is a notable opponent of countable additivity.

Kolmogorov then defines the *conditional probability of a given b* by the ratio of unconditional probabilities:

$$P(a | b) = \frac{P(a \cap b)}{P(b)}, \text{ provided } P(b) > 0.$$

Note that this ratio is undefined if either or both of the unconditional probabilities are undefined, or if $P(b) = 0$. Yet in uncountable spaces there can be genuine, non-trivial events whose probabilities are undefined (so-called ‘non-measurable sets’), and others whose probabilities are 0 (‘probability 0 does not imply impossible’ as textbooks, and Kolmogorov himself, caution us). So Kolmogorov’s definition does not guarantee that certain intuitive constraints on conditional probability are met—for example, that the probability of an event, *given itself*, is 1.

Kolmogorov addresses the probability 0 problem with a more sophisticated account of conditional probability as a random variable conditional on a sigma algebra, appealing to the Radon-Nikodym theorem to guarantee the existence of such a random variable. (See, e.g., Billingsley [1995].) A rival approach takes conditional probability $P(_, _)$ as primitive and defines the unconditional probability of a as $P(a, \mathbf{T})$, where \mathbf{T} is a necessary (e.g., tautological) proposition. Various axiomatizations of primitive conditional probability have been defended in the literature, typically differing only on the handling of conditional probabilities with zero unconditional probability antecedents. In many ways, the most general and elegant of the proposed axiomatizations is Popper's (1959). See Roeper and Leblanc (1999) for an encyclopedic discussion of competing theories of conditional probability, and Keynes (1921), Carnap (1950), Popper (1959), and Hájek (2003) for arguments that probability is inherently a two-place function.

Versions of *Bayes' theorem* can now be proven ([see BAYESIANISM](#)):

$$\begin{aligned} P(a | b) &= \frac{P(b | a)P(a)}{P(b)} \\ &= \frac{P(b | a)P(a)}{P(b | a)P(a) + P(b | \neg a)P(\neg a)} \end{aligned}$$

More generally, suppose there is a partition of hypotheses $\{h_1, h_2, \dots, h_n\}$, and evidence e . Then for each i ,

$$P(h_i | e) = \frac{P(e | h_i)P(h_i)}{\sum_{j=1}^n P(e | h_j)P(h_j)}.$$

The $P(e | h_i)$ terms are called *likelihoods*, and the $P(h_i)$ terms are called *priors*.

Finally, Kolmogorov defines a and b to be *independent* iff $P(a | b) = P(a)$; equivalently, iff $P(b | a) = P(b)$; equivalently, iff $P(a \cap b) = P(a)P(b)$ (for $P(a) \neq 0 \neq P(b)$). The terminology

suppresses the fact that such independence is really a *three*-place relation between an event, another event, and a *probability function*. This distinguishes probabilistic independence from such two-place relations as logical, causal, and counterfactual independence.

The next section turns to the so-called *interpretations* of probability: attempts to answer the central philosophical question: What is probability?

Frequentism

Ask a scientist what probability is, and one will typically get a *frequentist* answer: The probability of an event is the relative frequency of trials of a repeatable experiment on which that event occurs; sometimes the words ‘in the long run’ are added. This leaves open important questions: Which are the trials to be counted? How long does the run have to be? One may confine one’s attention to *actual* trials, realized in this world, or one may countenance *hypothetical* trials. And one may have merely finitely many trials to contend with, or one may have infinitely many, in which case probability will be identified with the *limit* of the relative frequency in a *sequence* of trials. One may thus immediately distinguish $2 \times 2 = 4$ variants of frequentism. However, the actual world typically delivers only finitely many trials of any given experiment. And it is often thought that if one is going to allow the trials to be hypothetical anyway, there is no obstacle to letting the sequence of trials be infinite, thus *guaranteeing* a ‘long run’. So one may confine one’s attention, as frequentists typically do, to just two of the possible positions: finite actual frequentism and infinite hypothetical frequentism.

In his discussion of the proportion of births of males and females, Venn contends that “probability is nothing but that proportion” (1866, 84, his emphasis)—a version of finite actual frequentism. Von Mises, by contrast, insists that probabilities exist only relative to virtual infinite sequences of ‘attributes’ called collectives. In a collective, the limiting relative frequency of any

attribute exists and is the same on any recursively specified subsequence. (Von Mises' original definition, in terms of "place selections", is here finessed by Church.)

The probability of a given attribute, relative to a collective, is then identified with its limiting relative frequency in that collective. Von Mises' position is thus a version of infinite hypothetical frequentism, as are those of Reichenbach and van Fraassen.

Any version of frequentism faces the notorious *reference class problem*. Any event, in all its detail, occurs exactly once, so if non-trivial frequencies are to be associated with it, it must be regarded as a token of a more general event type, whose instances constitute its reference class. However, there are indefinitely many ways of typing a given event. This would not be a problem if its relative frequency was the same in each reference class, or if one such class stood out as natural or privileged. The problem gains teeth to the extent that various competing reference classes have equal claim to determining the probability and that they yield different relative frequencies for the event.

In some cases the reference class problem may be solved for the actual, finite frequentist, but at the price of creating the equally notorious *problem of the single case*: intuitively, the objective probability of a one-off event may be less than 1, but finite frequentism cannot respect this intuition. Many events occur only once by any reasonable standard of typing: the 2000 presidential election, the invasion of Iraq, the last Lakers vs. Bulls game, and so on. The only natural reference class for such an event is the singleton set consisting of itself, and thus it has relative frequency 1 (and its non-occurrence has relative frequency 0). Nonetheless, it seems natural to think of non-extreme probabilities attaching to at least some of these 'single-case' events.

The problem of the single case is particularly striking, but there is really a sequence of related ‘granularity’ problems: the problem of the double case, the problem of the triple case ... A finite reference class of size n can only produce relative frequencies at a certain level of ‘grain’, namely $\frac{1}{n}$. Among other things, this rules out irrational probabilities; yet the best physical theories say otherwise (for example, various decay probabilities delivered by quantum mechanics are irrational). Furthermore, there is a sense in which any of these problems can be transformed into the problem of the single case. Suppose that a coin is tossed a thousand times. This can be regarded as a *single* trial of a thousand-tosses-of-the-coin experiment. Yet one does not want to be committed to saying that *that* experiment yields its actual result with probability 1.

The move to infinite hypothetical frequentism only makes the reference class problem worse. For not only must one choose a set of events in which to place a given event; since the set is now infinite, one must also choose an *ordering* among the events. After all, in non-trivial cases one can make the limiting relative frequency whatever value one likes simply by reordering the results of a given sequence. Consider the limiting relative frequency of even numbers among positive integers. On the ‘natural’ ordering $\langle 1, 2, 3, \dots \rangle$ it is $\frac{1}{2}$; however, one can make it $\frac{1}{4}$ by reordering the integers so that the even numbers occur at every fourth place in the sequence: $\langle 1, 3, 5, 2, 7, 9, 11, 4, 13, \dots \rangle$; and so on. Thus limiting relative frequencies are sensitive to apparently arbitrary choices of ordering, while it appears that probabilities are not. One might call this the *reference sequence problem*.

A sequence of events is said to be *exchangeable* with respect to a given probability function if all the joint probabilities of the events are invariant under finitely many permutations of the sequence: every event has the same probability, every conjunction of two events has the same

probability, every conjunction of three events has the same probability, and so on. A sequence of events is automatically exchangeable with respect to the relative frequency function: the frequency of an event is insensitive to *which* trials the event occurs at. Yet various events intuitively are not exchangeable with respect to the relevant probability function. Consider someone learning to throw a dart at a bull's eye: the sequence <MISS, MISS, HIT> is presumably more probable than <HIT, MISS, MISS>, because the dart-thrower's accuracy improves with practice. Yet the (finite) relative frequency of 'HIT' is $\frac{1}{3}$ either way. Since relative frequencies *force* a kind of symmetry that probabilities need not obey, they cannot be the same thing. (Ironically, it was the failure of a more thoroughgoing 'infinite exchangeability' that proved to be the undoing of hypothetical infinite frequentism in the previous paragraph.)

The classical interpretation

The brainchild of such founding fathers of probability as Pascal, Fermat, Huygens and Leibniz, and clearly articulated in Laplace (1814), the classical interpretation is the oldest interpretation of probability—indeed, it dates back to a time when the axiomatization and interpretation of probability were not clearly distinguished. It seeks to characterize the probability assignment of a rational agent in a state of epistemic neutrality with respect to a finite set of 'equipossibilities': the agent has either *no evidence*, or *symmetrically balanced evidence* regarding the possibilities. It appeals to the so-called *principle of indifference*: whenever there is no evidence favoring one possibility over another, each should be assigned the same probability as the others. So

$$P(e) = \frac{\text{number of equipossibilities in which } e \text{ occurs}}{\text{total number of equipossibilities}}$$

But the notion of ‘equipossibilities’ seems to presuppose some prior notion of probability. After all, the most obvious characterization of ‘symmetrically balanced evidence’ is in terms of equality of conditional probabilities: given evidence e and possible outcomes o_1, o_2, \dots, o_n , the evidence is symmetrically balanced with respect to the outcomes iff $P(o_1 | e) = P(o_2 | e) = \dots = P(o_n | e)$. Perhaps, then, one should regard the classical interpretation as an attempt to reduce quantitative probability to comparative probability: all numerical probabilities are ultimately based on facts about *equalities* among probabilities.

Note the structural resemblance of the classical theory to finite frequentism. Both theories see probability as a matter of even-handed counting and ratio-taking:

$$P(e) = \frac{\text{number of cases favorable to } e}{\text{total number of cases}}$$

It is just that for frequentism, the cases are *actual* outcomes of a *repeated* experiment, whereas for the classical theory they are *possible* outcomes of a *single* experiment. And indeed the classical theory faces many of the same problems as frequentism. There is the granularity problem: clearly, every classical probability is some fraction of the form $\frac{m}{n}$, where n is the number of possibilities. There is the exchangeability problem: classical probabilities are invariant under permutation of the labeling of the possibilities (for example, relabeling the faces of a die makes no difference to their probabilities of coming up). Thus, the classical interpretation cannot readily provide *asymmetric* probability distributions (e.g., for biased dice or coins), and it cannot handle distributions that evolve over time (e.g., for the dart-thrower’s hitting the bull’s eye).

Moreover, the reference class problem reappears. If one is truly ignorant about the results of some experiment, then presumably there is nothing to favor various competing choices of sample space. One should then be indifferent between, for example, {heads, tails} and {heads, tails,

edge}. And one should be indifferent between various refinements of the original space: for example, between spaces that refine in different ways the ‘heads’ outcome according to its final orientation relative to due north. Thus, probabilities will be determined by an apparently arbitrary choice of sample space. To adapt an example from physics, Bose-Einstein statistics, Fermi-Dirac statistics, and Maxwell-Boltzmann statistics each arise by considering the ways in which particles can be assigned to states, and then partitioning the set of alternatives in different ways (see STATISTICAL MECHANICS). (See, e.g., Fine [1973].) Someone ignorant of which statistics apply to a given type of particle can only make an arbitrary choice and hope for the best.

In typical applications of the classical theory—gambling games, for example—one is not wholly ignorant, but the evidence that one has is symmetrically balanced regarding the possibilities. There are two problems here: in the ‘evidence’, and in the ‘symmetry’. Classical probabilities are acutely sensitive to the evidence. If the evidence becomes *unbalanced*, favoring some outcomes over others, then classical probabilities are not merely revised, they are *destroyed*. And there may be competing respects of symmetry, each equally compelling. This problem arises especially when there are infinitely many possible outcomes. Then, the equipossibilities must be a finite partition of the outcomes. But *which* partition?

A tempting answer may be: the most “natural” partition. However, ‘Bertrand’s paradoxes’ show that there need not be any such. The trick is to give competing parametrizations of a given problem that are non-linearly related to one another, but equally “natural”. Suppose one is told only that a car traveled 100 miles at an average speed between 50 and 100 m.p.h. What is the probability that its average speed was between 75 and 100 m.p.h? Perhaps: 0.5—since (50, 75) and (75, 100) are equipossible intervals for the average speed. But the question could be

equivalently formulated: a car took between 1 and 2 hours to travel 100 miles. What is the probability that it took between 1 hour and $1\frac{1}{3}$ hours? Now it seems that there are three equipossible intervals for the time taken: $(1, 1\frac{1}{3})$, $[1\frac{1}{3}, 1\frac{2}{3})$, and $[1\frac{2}{3}, 2)$; whence the answer should be $\frac{1}{3}$.

Logical probability

Many philosophers—Leibniz, von Kries, Keynes, Wittgenstein, Waismann, Carnap, and others—have tried to explicate the following “logical” concept of conditional probability:

$$P(p | q) = \frac{\text{the proportion of logically possible worlds in which both } p \text{ and } q \text{ are true}}{\text{the proportion of logically possible worlds in which } q \text{ is true}}$$

An obvious problem has been to justify a *measure* of the “proportion of logically possible worlds in which a proposition is true”. Early attempts—including those by Carnap that will be the focus here—tried to apply the controversial principle of indifference (see RUDOLF CARNAP). Carnap’s early (1950) constructions are very similar to systems developed earlier by W.E. Johnson (1921). See (INDUCTIVE LOGIC for further references on Carnapian inductive logic and logical probability).

Begin with a first-order language L containing a finite number of monadic predicates: F , G , H , ..., and a finite or denumerable number of individual constants a , b , c , Then define an (“*a priori*”) unconditional probability function $P(\bullet)$ over the sentences of L , in a way that only appeals to their *syntactic structure* (whence the name “logical” probability). Finally, use the standard ratio definition to construct a conditional probability function $P(\bullet | \bullet)$ over pairs of sentences of L .

The results of this procedure will be *language-relative*: if one describes the same phenomena by means of a different language L^* —equipped with a different stock of monadic predicates—one will typically not recover the same probabilities. Consider two languages used to represent the outcomes of random draws from an urn filled with colored balls. Let L contain the color predicates “blue” and “green”, L^* the predicates “grue” and “bleen”. The intended interpretation: a draw is grue just in case it is one of the first million and green or a later one and blue; a draw is bleen just in case it is one of the first million and blue or a later one and green. Starting with L , use whatever is the appropriate procedure to calculate $P(\text{draw 1,000,001 is green} \mid \text{the first 1,000,000 draws are green})$. Starting with L^* , use this procedure to calculate $P(\text{draw 1,000,001 is grue} \mid \text{the first 1,000,000 draws are grue})$. If syntax is all that matters, then these conditional probability values will be identical—and surely greater than $\frac{1}{2}$, at least if logical probability is to have a hope of modeling actual inductive reasoning (see INDUCTIVE LOGIC). The trouble is that the second conditional probability, translated into L , is just $P(\text{draw 1,000,001 is blue} \mid \text{the first 1,000,000 draws are green})$. One can avoid contradiction, but only by explicitly insisting that probability is language-relative. And that raises a serious problem—really, the reference class problem in a new guise: if one wishes to employ logical probability as a foundation for inductive inference, which is the “right” language to use? The remainder of this discussion will presuppose that an answer to this question has been found (for Carnap, this question was “external” to inductive logic anyway, and his later systems did not have this blatant form of language relativity—see Carnap [1980] for discussion).

Returning now to Carnap’s early systems, consider a simple language with only two monadic predicates ‘ F ’ and ‘ G ’ and only two individual constants ‘ a ’ and ‘ b ’. This language yields exactly 16 maximally specific descriptions of the world—the *state descriptions* of L : ($Fa \wedge Ga$

$\wedge Fb \wedge Gb$, $(Fa \wedge Ga \wedge Fb \wedge \neg Gb)$, etc. Two state descriptions S_1 and S_2 are *permutations* of each other if S_1 can be obtained from S_2 by some permutation of the individual constants. For example, $Fa \wedge \neg Ga \wedge \neg Fb \wedge Gb$ and $\neg Fa \wedge Ga \wedge Fb \wedge \neg Gb$ are permutations of each other. A *structure description* in L is a disjunction of state descriptions, closed under permutation. L provides these 10 structure descriptions:

$$\begin{array}{ll}
Fa \wedge Ga \wedge Fb \wedge Gb & (Fa \wedge \neg Ga \wedge \neg Fb \wedge Gb) \vee (\neg Fa \wedge Ga \wedge Fb \wedge \neg Gb) \\
(Fa \wedge Ga \wedge Fb \wedge \neg Gb) \vee (Fa \wedge \neg Ga \wedge Fb \wedge Gb) & (Fa \wedge \neg Ga \wedge \neg Fb \wedge \neg Gb) \vee (\neg Fa \wedge \neg Ga \wedge Fb \wedge \neg Gb) \\
(Fa \wedge Ga \wedge \neg Fb \wedge Gb) \vee (\neg Fa \wedge Ga \wedge Fb \wedge Gb) & \neg Fa \wedge Ga \wedge \neg Fb \wedge Gb \\
(Fa \wedge Ga \wedge \neg Fb \wedge \neg Gb) \vee (\neg Fa \wedge \neg Ga \wedge Fb \wedge Gb) & (\neg Fa \wedge Ga \wedge \neg Fb \wedge \neg Gb) \vee (\neg Fa \wedge \neg Ga \wedge \neg Fb \wedge Gb) \\
Fa \wedge \neg Ga \wedge Fb \wedge \neg Gb & \neg Fa \wedge \neg Ga \wedge \neg Fb \wedge \neg Gb
\end{array}$$

Now, assign non-negative real numbers to the state descriptions, so that these 16 numbers sum to one. Any such assignment will constitute an (“*a priori*”) unconditional probability function $P(\bullet)$ over the state descriptions of L . To extend $P(\bullet)$ to the entire language L , note that the probability of a disjunction of mutually exclusive sentences is the sum of the probabilities of its disjuncts. Since every sentence in L is equivalent to some disjunction of state descriptions, and all the state descriptions are mutually exclusive, this gives a complete unconditional probability function $P(\bullet)$ over L —typically called a *measure function*. The standard ratio definition then yields a conditional probability function $P(\bullet \mid \bullet)$ over pairs of sentences in L .

Carnap (1950) discusses two “natural” measure functions. The first, m^\dagger , treats each state description as *equiprobable a priori*: if there are N state descriptions in L , then m^\dagger assigns $\frac{1}{N}$ to each. However natural this measure function may seem, it has the consequence that the resulting probabilities cannot undergird *learning from experience*. To see why, observe that

$$P(Fb \mid Fa) = \frac{m^\dagger(Fb \wedge Fa)}{m^\dagger(Fa)} = \frac{1}{2} = m^\dagger(Fb) = P(Fb).$$

So ‘learning’ that one object has property F cannot affect the probability that any other object will also have property F . Indeed, it can be shown that *no matter how many objects are assumed to be F* , this will *always be irrelevant* (according to probability functions based on m^\dagger) to the hypothesis that a distinct object will also be F —a feature widely viewed as a serious shortcoming of m^\dagger .

As a result, Carnap formulated an alternative measure function m^* : First, assign equal probabilities to each *structure* description. Then, each state description entailing a given structure description is assigned an equal portion of the probability assigned to that structure description. So, in the present toy language, the state description ‘ $Fa \wedge Ga \wedge \neg Fb \wedge Gb$ ’ gets assigned *a priori* probability of $\frac{1}{20}$ ($\frac{1}{2}$ of $\frac{1}{10}$), but the state description ‘ $Fa \wedge Ga \wedge Fb \wedge Gb$ ’ receives an *a priori* probability of $\frac{1}{10}$ ($\frac{1}{1}$ of $\frac{1}{10}$). Unlike m^\dagger , m^* *does* allow for “learning from experience”: e.g. $P(Fa \mid Fb) = \frac{3}{5} > \frac{1}{2} = P(Fa)$. Still, even m^* can give unintuitive results in more complex languages (see Carnap [1952] for discussion). Also, note that the state descriptions are exchangeable with respect to m^* , an omen that logical probabilities will face some of the problems that plagued the frequentist and the classical probabilist.

Carnap (1952) presents a more complicated “continuum” of conditional probability functions. This continuum depends on a parameter λ intended to reflect the “speed” with which learning from experience is possible. $\lambda = 0$ corresponds to the “straight rule”, which says that the probability that the next object observed will be F , conditional upon a sequence of past

observations, is simply the frequency of F objects in that sequence; $\lambda = +\infty$ yields a conditional probability function much like that derived from the measure function m^* (*i.e.*, $\lambda = +\infty$ implies that there is no learning from experience); $\lambda = \kappa$ (which is the number of independent families of predicates in Carnap's more elaborate [1952] linguistic framework) yields a conditional probability function equivalent to that generated by the measure function m^* .

Problems remain. None of the Carnapian systems allow universal generalizations to have non-zero probability. (Hintikka, and Hintikka and Niiniluoto provide alterations of the Carnapian framework that *do* allow for non-zero probabilities of universal generalizations.) Carnap's early systems also failed to allow for *analogical effects*, since according to them the fact that two objects share several properties in common is (in many cases) *irrelevant* to whether they share any *other* properties in common. Carnap's most recent (and most complex) theories of logical probability (1980) include two additional parameters designed to provide the theory with enough flexibility to overcome these (and other) limitations. Unfortunately, no Carnapian logical theory of probability to date has dealt successfully with the problem of analogical effects (see Maher 2001 for further discussion). The consensus now seems to be that the Carnapian project of constructing an adequate logical theory of probability is all but hopeless: the syntactical constraints implicit in any such theory will inevitably prevent the theory from being able to model certain essential features of statistical inference and/or inductive logic (see INDUCTIVE LOGIC).

Subjectivism

In slogan form, subjectivism regards probabilities as *degrees of belief*, or *credences*. But what are credences? Subjectivists since Ramsey (1926) have insisted that they must be intimately

tied to the behavioral dispositions of suitable agents. On one influential account, advocated by de Finetti (1937),

an agent's credence in e is p

iff

p units of utility is the price at which the agent would buy or sell a bet that pays 1 unit of utility if e , 0 if $\neg e$.

This is at best a first approximation to an analysis of credence. One surely should allow the buying and selling prices of at least some bets to come apart. And even when they agree, there are problems. How does one separate the agent's epistemic attitude *to e* from his or her attitude (favorable, unfavorable, or neutral) to gambling? Indeed, one may insist on separating epistemic attitudes from desire-based attitudes altogether; one can imagine, for example, a chronic apathetic who has opinions, but who lacks corresponding desires (for bets, or for anything). Moreover, the very placement of the bet may change the world in ways that affect the agent's credences.

Be that as it may, there are famous arguments that credences must conform to the probability calculus, at least if one demands that the agent be in some sense *ideally rational*. For example, if one's credences do *not* so conform, one is susceptible to a *Dutch Book*, a sequence of bets that one regards as acceptable taken individually, but that collectively guarantee one's loss, however the world turns out. Conversely, if one's credences *do* so conform, one is immune to a Dutch Book. Rationality, it is concluded, requires obedience to the probability calculus ([see DUTCH BOOK ARGUMENT](#)).

Utilities (desirabilities) of outcomes, their probabilities, and rational preferences are all intimately linked. *The Port Royal Logic* showed how utilities and probabilities together determine rational preferences; de Finetti's betting interpretation derives probabilities from utilities and rational preferences; von Neumann and Morgenstern (1944) derive utilities from probabilities and rational preferences. And most remarkably, Ramsey (1926) (and later, Savage

[1954] and Jeffrey [1966]) derives *both* probabilities *and* utilities from rational preferences alone. (See RAMSEY, FRANK PLUMPTON.)

First, Ramsey defines a proposition to be *ethically neutral*—relative to an agent and an outcome—if the agent is indifferent between having that outcome when the proposition is true and when it is false. Suppose that the agent prefers a to b . Then an ethically neutral proposition n has probability $\frac{1}{2}$ iff the agent is indifferent between the gambles:

a if n , b if not

b if n , a if not.

One may assign arbitrarily to a and b any two real numbers $u(a)$ and $u(b)$ such that $u(a) > u(b)$, thought of as their respective desirabilities. Having done this for the one arbitrarily chosen pair a and b , the utilities of all other propositions are determined. Given various assumptions about the richness of the preference space, and certain 'consistency assumptions', Ramsey can define a real-valued utility function of the outcomes a , b , etc—in fact, various such functions will represent the agent's preferences. He is then able to define equality of differences in utility for any outcomes over which the agent has preferences. It turns out that ratios of utility-differences are invariant—the same whichever representative utility function one chooses. This fact allows Ramsey to define degrees of belief as ratios of such differences. For example, suppose the agent is indifferent between a , and the gamble " b if x , c otherwise". Then his or her degree of belief in x , $P(x)$, is given by:

$$P(x) = \frac{u(a) - u(c)}{u(b) - u(c)}.$$

Ramsey shows that degrees of belief so derived obey the probability calculus (with finite additivity). He calls what results "the logic of partial belief".

Ramsey avoids some of the objections to the betting interpretation, but not all of them. Notably, the essential appeal to gambles again raises the concern that the wrong quantities are

being measured. And his account has new difficulties. It is unclear what facts about agents fix their preference rankings. It is also dubious that *consistency* alone requires one to have a set of preferences as rich as Ramsey requires, or that one can find ethically neutral propositions of probability $\frac{1}{2}$. This in turn casts some doubt on Ramsey's claim to assimilate probability theory to logic.

Savage (1954) likewise derives probabilities and utilities from preferences among options that are constrained by certain putative 'consistency' principles. For a given set of such preferences, he generates a class of utility functions, each a positive linear transformation of the other (i.e. of the form $u_1 = au_2 + b$, where $a > 0$), and a unique probability function. Together these are said to 'represent' the agent's preferences. Jeffrey (1966) refines the method further. The result is theory of decision according to which rational choice maximizes 'expected utility', a certain probability-weighted average of utilities.

So far, this is a static picture of a rational agent. How should one update one's degrees of belief in the light of new evidence? The favored rule among subjectivists is *conditionalization*: where e is the strongest proposition of which one becomes certain, one's new credence function is related to the old by:

$$(\text{Conditionalization}) \quad C_{\text{new}}(\bullet) = C_{\text{old}}(\bullet \mid e),$$

using here and in what follows " $C(\bullet)$ " to distinguish credence from other kinds of probability.

So-called *subjective Bayesianism* holds that an agent's epistemic trajectory is rational iff at any moment his her credences are representable by a probability function, and he or she always updates by conditionalization. This is at once a highly demanding and highly permissive epistemology. It is demanding, because conformity to probability theory is demanding. It is permissive, because there is no requirement that degrees of belief in any way correspond to the

way the world is. So someone who assigns probability 1 to the universe being ruled by a rubber chicken can meet the Bayesian standards for rationality—as long as they obey the probability calculus in all their other assignments and always update by conditionalizing. Bayesians reply that various convergence theorems show roughly that in the long run, agents who do not give probability 0 to genuine possibilities, and whose stream of evidence is sufficiently rich, will eventually be arbitrarily close to certain regarding the truth about the world in which they live. For skepticism about the value of these theorems, see Earman (1992).

In any case, there are numerous proposals for further constraints on priors. Some—e.g., by Jeffreys and Jaynes—appeal to a version of the principle of indifference. Some can be regarded as instances of a certain schema, proposed by Gaifman (1988). He coins the term “expert probability” for a probability assignment that a given agent strives to track, codifying this idea as follows:

$$(\text{Expert}) \quad C(a \mid pr(a) = x) = x, \text{ for all } x \text{ such that } C(pr(a) = x) > 0.$$

Here $pr(a)$ is the assignment that the agent regards as expert. For example, if one regards the local weather forecaster as an expert, and he or she assigns probability 0.1 to it raining tomorrow, then one may well follow suit:

$$C(\text{rain} \mid pr(\text{rain}) = 0.1) = 0.1.$$

More generally, one might speak of an entire probability function as being such a guide for an agent, over a specified set of propositions—so that (Expert) holds for any choice of A from that set. A *universal expert function* would guide *all* of the agent’s probability assignments in this way. Van Fraassen (1995) argues that an agent’s *future* probability functions are universal expert functions for that agent—his *Reflection Principle*:

$$C_t(a \mid C_t(a) = x) = x, \text{ for all } a \text{ and for all } x \text{ such that } C_t(C_t(a) = x) > 0,$$

where C_t is the agent's probability function at time t , and $C_{t'}$ his or her function at later time t' .

The principle encapsulates a certain demand for ‘diachronic coherence’ imposed by rationality.

Van Fraassen defends it with a ‘diachronic’ Dutch Book argument (one that considers bets placed at different times), and by analogizing violations of it to the sort of pragmatic inconsistency that one finds in Moore’s paradox. For example, suppose that one is certain that

one will tomorrow assign probability $\frac{1}{2}$ to it raining the day after, but that one nonetheless

assigns it probability $\frac{1}{3}$ now. While this is not logically inconsistent, it is surely puzzling.

One may go still further. There may be universal expert functions for all rational agents. The *Principle of Direct Probability* regards the relative frequency function as a universal expert function. Let a be an event-type, and let $relfreq(a)$ be the relative frequency of a (in some suitable reference class). Then for any rational agent, one has:

$$C(a \mid relfreq(a) = x) = x, \text{ for all } a \text{ and for all } x \text{ such that } C(relfreq(a) = x) > 0.$$

The next section takes up what many consider the most important such universal expert function.

Objective Chance

De Finetti, the great probabilist, quipped that “Probability does not exist” (1990, 1). What he meant was that *all* probability is *subjective*. Yet there is a strong *prima facie* case for recognizing the existence of *objective chances*: probabilities that attach to physical systems and their behavior independently of anyone’s mental state, and that capture *contingent* facts about those systems, and not merely quasi-logical relations among propositions concerning them. One wonders, for example, whether a certain coin is biased, and if so, to what degree and in what

direction. Translation: One wonders what the *chance of heads* would be, were the coin tossed fairly.

This example would not faze a committed subjectivist such as de Finetti, nor a frequentist like von Mises (1957) who denies that probability ever applies in the single case. But more serious examples from physics suggest that, ultimately, resistance is futile. For starters, *statistical physics* says that the entire universe *could* evolve towards a state of lower entropy, but that the chance of its doing so is vanishingly small. Such chances are seemingly compatible with determinism at the level of the fundamental dynamical laws (this is controversial: see IRREVERSIBILITY; STATISTICAL MECHANICS), and following Bernoulli, various authors have for that reason doubted their credentials. However, see Levi (1990) for a valuable discussion of authors since Cournot and Venn who countenance such compatibilist chances. The compatibility is putatively secured by relativizing chances to a kind of trial. For example, a coin may have chance $\frac{1}{2}$ of landing heads relative to the specification of being tossed high above a flat surface, say, while having a chance 1 of landing heads relative to a precise specification of the initial conditions of a particular toss in a deterministic world. Levi argues that the applicability of chance hypotheses and statistical techniques does not presuppose an underlying indeterminism, and so a theory of chance should remain neutral vis-à-vis determinism.

In any case, commitment to chances has a second source: “*collapse*” theories of quantum mechanics explicitly introduce indeterministic dynamical laws that not only specify what courses of evolution are possible for a given physical system with a given initial state, but also specify exact *probabilities* for each such trajectory. (See QUANTUM MEASUREMENT PROBLEM; QUANTUM MECHANICS.) The subjectivist or von Mises-style frequentist seems left only with the option of denying—from the armchair!—that the physical theories that postulate them are

true or coherent. It is better simply to acknowledge that objective chances are or at least could be real, and then to go on to consider what sort of account one could give of them.

Reductionist accounts attempt to reduce facts about objective chances to the totality of non-modal facts about a world. *Non-reductionist* accounts deny that chance even supervenes on the non-modal facts. Actual frequentism is clearly a reductionist view. More sophisticated is Lewis's "best systems" approach (1994), which sees the laws for a world w as, roughly, being theorems of that axiomatic system for describing non-modal facts about w that achieves an optimal balance of simplicity and informativeness. In the case of probabilistic laws, Lewis invokes a third criterion: a system for w is 'better' to the extent that it assigns a higher probability to the total history of w . The simplest form of non-reductionism is *primitivism*, which takes chances to be unanalyzable features of the world. Alternatively, one might try to explain one's non-reductionist chances by appeal to some other bit of metaphysical gadgetry—e.g. Armstrong takes chances to consist in higher-order relations of "probabilification" that obtain between universals.

No mention has been made so far of "propensity" interpretations of probability. Everyone, reductionist and non-reductionist alike, can agree that in a chancy world, some physical systems will have "propensities" to exhibit certain behaviors under certain conditions. For all one need *mean* by that is that counterfactuals of the following form are true of these systems: "were the system in conditions C , there would be a chance of x that it would manifest behavior B ". So propensities—understood as tendencies, or variable-strength dispositions—can be analyzed straightforwardly in terms of subjunctive conditionals whose consequents make reference to objective chances.

Propensity *interpretations* of probability aim to reverse this order of analysis, explaining objective chances directly in terms of propensities. For some authors (Popper, Gillies), chances

are dispositions for a chance set-up to produce long-run relative frequencies; for others (Giere, Fetzer, Miller) they are dispositions for a chance set-up to produce outcomes on single trials. Subtle variations can be found in the work of Hacking, Mellor, and Levi. (See Gillies 2000 and Hájek 2003 for surveys.)

But there are some general problems that any propensity account faces. Suppose that system S has a certain tendency to manifest behavior B under conditions C . One must be able to attach numbers to such a tendency as a measure of its strength *without* appealing to the concept of chance; it is not clear how this is to be done, nor why the results should obey the probability calculus. Moreover, how do propensities for distinct systems yield propensities for the composite systems they make up? Here are two coins, each with a propensity of 0.5 of landing heads if tossed. Suppose both are tossed at once. If there is a chance that both will land heads, then there must be a propensity possessed by the *combined* two-coin system. If so, what guarantees that the marginal probabilities (for each coin considered separately) will be recovered correctly from this composite propensity? And one cannot stop here: one had better say, of *the world as a whole*, that it exhibits, at each moment, propensities to evolve in various different ways. Having gone so far, one might as well simply say that instead of exhibiting “propensities”, the world exhibits *chances*, thus avoiding (by stipulation) the original problem of their conformity to the probability calculus—and thus arriving at primitivism about chance. If that is right, then it is not clear that propensity accounts offer a genuinely new option for understanding probability.

Although distinct, objective and subjective probability display an extremely important connection. Lewis (1980) formulates it in his Principal Principle:

$$(PP) \quad C_o(a \mid e \wedge ch_f(a) = x) = x.$$

Here C_0 is some reasonable “initial” (*a priori*) credence function, a an arbitrary proposition, $ch_t(a) = x$ the claim that the chance, at time t , of a is x , and e an “admissible” proposition—one that does not contain information relevant to a beyond that given by its chance at t . (Thus, e.g., a itself is inadmissible.)

One can apply (PP) to a “non-initial” agent by modeling her credence C as the result of conditionalizing some reasonable initial credence C_0 on some suitable evidence. Let h describe a complete possible course of history until time t . Let l describe some possible fundamental laws compatible with h , and assume that the way in which chances depend on history is underwritten by these laws. Then the conjunction $h \wedge l$ picks out a unique chance-distribution $P(\bullet)$ for time t . Thus, if a proposition of the form $(h \wedge l \wedge ch_t(a) = x)$ is consistent, then the third conjunct is entailed by the first two. Assuming, as seems reasonable, that the conjunction $h \wedge l$ is admissible, it follows that

$$(\text{PP}^*) \quad C_0(a \mid h \wedge l) = P(a).$$

Much of the debate between reductionists and non-reductionists consists in a war of intuitions: e.g., the reductionist claims to find the non-reductionist’s extra, irreducibly modal feature of metaphysical reality unintelligible; the non-reductionist claims to “show” that distinct chances can give rise to exactly the same total histories of non-modal fact—a draw, perhaps. But (PP) and (PP^{*}) appear to open up new lines of argument.

The non-reductionist alleges that reductionism is *inconsistent* with (PP^{*}). Typical reductionist views will allow that the chance-laws can have some non-zero chance of failing to obtain. For the reductionist says that these laws are determined by the total history of non-modal fact. But these laws issue in chance-distributions over possible total histories of non-modal fact. Thus, it may turn out that positive chances are assigned to total histories that would specify

different laws—the so-called “undermining” of the chance-laws by themselves. (See Lewis [1994].) Example: A coin is about to be tossed exactly 10^{10} times. As it happens, exactly half the tosses will land heads. A reductionist might say that it *follows* that the chance of heads on each toss is 0.5, adding that the correct chance-laws will treat the tosses as independent. So there is now a large chance that the frequency of heads will be *different* from what it actually is—and if so, the *laws* will be different as well.

The inconsistency with (PP*) is now manifest. Consider those consistent history-law conjunctions $h \wedge l$ that entail that $P(l) < 1$. Pick such a conjunction; by (PP*),

$$C_0(l \mid h \wedge l) = P(l) < 1.$$

But by the probability calculus,

$$C_0(l \mid h \wedge l) = 1.$$

Lewis responds by amending (PP*) to what he calls the “New Principle”:

$$(NP) \quad C_0(a \mid h \wedge l) = P(a \mid l),$$

thus avoiding the inconsistency. The consensus in the literature seems to be that unlike the original Principal Principle, (NP) is unintuitive and, in application, unwieldy.

The reductionist (e.g. Lewis [1994]) retorts that non-reductionists are hard-pressed to show how chances, understood their way, constrain rational credences according to (PP). But can the reductionist himself meet this challenge? Presumably, he ought to provide a *derivation* of (PP) from constraints on rational credence to which he is *already* committed, and his reductionist analysis of chance. The literature provides no such derivation. And while a non-reductionist may be unable to supply such a derivation it is not clear why she *needs* to. Arguably, *all* are committed to the existence of substantive constraints on rational credence; why can't the non-

reductionist simply include (PP) as one of them? (See Hall 2003 for further discussion.) Perhaps, then, the debate between reductionism and non-reductionism remains a stalemate.

Finally, a ‘deflationary’ account of chance, associated with de Finetti and his followers, has proved to be very influential. Consider an infinite exchangeable sequence of events with respect to a probability function P . *De Finetti’s representation theorem* states that the probability according to P of exactly k of the events occurring in n trials is given by

$$\int_0^1 \binom{n}{k} p^k (1-p)^{n-k} f(p) dp,$$

for all n and k and for some density function f . The upshot is that any such probability distribution is representable as a ‘weighted average’ of distributions. Each distribution corresponds to a hypothesis about the value of the probability p of an event occurring on a single trial; it gives the probability of k such events occurring in n independent, identically distributed trials, given that fixed value of p . One can then average these distributions using the probabilities of their corresponding hypotheses about the value of p as weights. The result is significant because it enables a subjectivist to ‘simulate’ an objectivist about chance when the exchangeability assumption holds, and for many situations this seems reasonable. If P is one’s subjective probability function, then it is *as if* one spread probability over various hypotheses about the single case objective chance of the event, which remains fixed across infinitely many independent trials of the experiment in question. See Skyrms (1994) for an excellent discussion of generalizations of exchangeability and their use in formulating various Goodmanian theses about projectability. Indeed, commonsense often (but not invariably) seems to require one’s probabilities to be exchangeable over ‘green’-like hypotheses, but not ‘grue’-like hypotheses.

Conclusion

Feller (1957, 19) writes: “All possible definitions of probability fall short of the actual practice.” Certainly, a lot is asked of the concept of probability. It is supposed at once to capture a quasi-logical notion, a subjective notion, and an objective notion instantiated in the mind-independent world. Perhaps one would do better to think of these as distinct *concepts* of probability. Each of the leading interpretations, then, attempts to illuminate one of these concepts, while leaving the others in the dark. In that sense, the interpretations might be regarded as complementary, although to be sure each may need some further refinement. Clearly, much work remains to be done on the philosophical foundations of probability. Equally clearly, we have come a long way since the *Port Royal Logic*.

References

Note: due to space limitations, not all of the references could be given for the authors cited in the text. Many further references can be found in Hájek (2003), and in **INDUCTIVE LOGIC**.

Arnauld, Antoine (1964), *Logic, or, The Art of Thinking* ("The Port Royal Logic"), (tr. J. Dickoff and P. James) Indianapolis: Bobbs-Merrill.

Billingsley, Patrick (1995), *Probability and Measure*, 3rd ed. New York: John Wiley & Sons.

Carnap, Rudolf (1950), *Logical Foundations of Probability*. Chicago: University of Chicago Press.

— (1952), *The Continuum of Inductive Methods*. Chicago: The University of Chicago Press.

—(1962), *Logical Foundations of Probability*, 2nd ed. Chicago: University of Chicago Press.

—(1980), “A Basic System of Inductive Logic II”, in Richard Jeffrey (ed.), *Studies in Inductive Logic and Probability, Volume II*. Berkeley: University of California Press, 7-155.

De Finetti, Bruno (1937), “La Prévision: Ses Lois Logiques, Ses Sources Subjectives”, *Annales de l'Institut Henri Poincaré*, 7: 1-68. Translated as “Foresight. Its Logical Laws, Its Subjective Sources”, in H. E. Kyburg, Jr. and H. E. Smokler (eds.), *Studies in Subjective Probability*. New York: Robert E. Krieger Publishing Company.

—(1990), *Theory of Probability*, Vol. 1. Chichester: Wiley Classics Library, John Wiley & Sons.

Earman, John (1992), *Bayes or Bust?*. Cambridge: MIT Press.

Feller, William (1957), *An Introduction to Probability Theory and Its Applications*. New York: John Wiley & Sons, Inc.

Fine, Terrence (1973), *Theories of Probability*. New York: Academic Press.

Gaifman, Haim (1988), “A Theory of Higher Order Probabilities”, in Brian Skyrms and William L. Harper (eds.), *Causation, Chance, and Credence*. Dordrecht: Kluwer Academic Publishers, .

Gillies, Donald (2000), “Varieties of Propensity”, *British Journal for the Philosophy of Science* 51: 807-835.

Hájek, Alan (2003), "Interpretations of Probability", in Edward N. Zalta (ed.), *The Stanford Encyclopedia of Philosophy* (Summer 2003 Edition), URL:
<http://plato.stanford.edu/archives/sum2003/entries/probability-interpret/>.

— (2003), "What Conditional Probability Could Not Be", *Synthese* 137: 273-323.

Hall, Ned (2003), “Two Mistakes About Credence and Chance”, *Australasian Journal of Philosophy*: 93-111.

Jeffrey, Richard (1983), *The Logic of Decision*, 2nd ed. Chicago: University of Chicago Press.

Johnson, William E. (1921), *Logic*. Cambridge: Cambridge University Press.

Keynes, John M. (1921), *A Treatise on Probability*. London: Macmillan.

Kolmogorov, Andrei. N. (1933), *Grundbegriffe der Wahrscheinlichkeitrechnung, Ergebnisse Der Mathematik*. Translated as *Foundations of Probability*. New York: Chelsea Publishing Company, 1950.

Laplace, Pierre Simon (1951), *A Philosophical Essay on Probabilities*, English edition. New York: Dover Publications Inc. (Original published 1814.)

Levi, Isaac (1990), “Chance”, *Philosophical Topics* 18: 117-148.

Lewis, David (1980), “A Subjectivist’s Guide to Objective Chance”, reprinted with postscripts in Lewis 1986: 83-132.

—(1986), *Philosophical Papers: Volume II*. New York: Oxford University Press.

—(1994), “Humean Supervenience Debugged”, *Mind* 103: 473-90.

Maher, Patrick (2001), “Probabilities for Multiple Properties: the Models of Hesse and Carnap and Kemeny”, *Erkenntnis* 55: 183-216.

Muckenheim, W., G. Ludwig, C. Dewdney, P. Holland, A. Kyriyanidis, J. Vigier, N. Petroni, M. Bartlett, and E. Jaynes (1986), “A Review of Extended Probability”, *Physics Reports* 133: 337–401.

Popper, Karl (1959), *The Logic of Scientific Discovery*. London: Hutchinson & Co.

Ramsey, Frank P. (1926), “Truth and Probability”, in R. B. Braithwaite (ed.), *Foundations of Mathematics and other Essays*. Routledge & P. Kegan. Reprinted in H. E. Kyburg, Jr. and H.

E. Smokler (eds.), *Studies in Subjective Probability*, 2nd ed., R. E. Krieger Publishing Company, 1980: 23-52.

Renyi, Alfred (1970), *Foundations of Probability*. Holden-Day, Inc.

Roeper, Peter, and Hughes Leblanc (1999), *Probability Theory and Probability Logic*. Toronto: University of Toronto Press.

Savage, Leonard J. (1954), *The Foundations of Statistics*. New York: John Wiley.

Skyrms, Brian (1980), *Causal Necessity*. New Haven: Yale University Press.

—(1994), “Bayesian Projectibility”, in D. Stalker (ed.) *Grue: Essays on the New Riddle of Induction*. Chicago: Open Court: 241-262.

van Fraassen, Bas (1995), “Belief and the Problem of Ulysses and the Sirens”, *Philosophical Studies* 77: 7-37.

Venn, John (1866), *The Logic of Chance*. London and Cambridge: Macmillan.

von Mises, Richard (1957), *Probability, Statistics and Truth*, revised English edition. New York: Macmillan.

von Neumann, John, and Oskar Morgenstern (1944), *Theory of Games and Economic Behavior*. Princeton: Princeton University Press.

Branden Fitelson, Alan Hájek, and Ned Hall