

Introduction to Time Series Analysis. Lecture 2.

Peter Bartlett

Last lecture:

1. Objectives of time series analysis.
2. Time series models.
3. Time series modelling: Chasing stationarity.

Introduction to Time Series Analysis. Lecture 2.

Peter Bartlett

1. Stationarity
2. Autocovariance, autocorrelation
3. MA, AR, linear processes

Stationarity

$\{X_t\}$ is **strictly stationary** if

for all $k, t_1, \dots, t_k, x_1, \dots, x_k$, and h ,

$$P(X_{t_1} \leq x_1, \dots, X_{t_k} \leq x_k) = P(X_{t_1+h} \leq x_1, \dots, X_{t_k+h} \leq x_k).$$

i.e., shifting the time axis does not affect the distribution.

We shall consider **second-order properties** only.

Mean and Autocovariance

Suppose that $\{X_t\}$ is a time series with $E[X_t^2] < \infty$.

Its **mean function** is

$$\mu_t = E[X_t].$$

Its **autocovariance function** is

$$\begin{aligned}\gamma_X(s, t) &= \text{Cov}(X_s, X_t) \\ &= E[(X_s - \mu_s)(X_t - \mu_t)].\end{aligned}$$

Weak Stationarity

We say that $\{X_t\}$ is **(weakly) stationary** if

1. μ_t is independent of t , and
2. For each h , $\gamma_X(t + h, t)$ is independent of t .

In that case, we write

$$\gamma_X(h) = \gamma_X(h, 0).$$

Stationarity

The autocorrelation function (ACF) of $\{X_t\}$ is defined as

$$\begin{aligned}\rho_X(h) &= \frac{\gamma_X(h)}{\gamma_X(0)} \\ &= \frac{\text{Cov}(X_{t+h}, X_t)}{\text{Cov}(X_t, X_t)} \\ &= \text{Corr}(X_{t+h}, X_t).\end{aligned}$$

Stationarity

Example: i.i.d. noise, $E[X_t] = 0$, $E[X_t^2] = \sigma^2$. We have

$$\gamma_X(t+h, t) = \begin{cases} \sigma^2 & \text{if } h = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Thus,

1. $\mu_t = 0$ is independent of t .
2. $\gamma_X(t+h, t) = \gamma_X(h, 0)$ for all t .

So $\{X_t\}$ is stationary.

Similarly for any white noise (uncorrelated, zero mean), $X_t \sim WN(0, \sigma^2)$.

Stationarity

Example: Random walk, $S_t = \sum_{i=1}^t X_i$ for i.i.d., mean zero $\{X_t\}$.
We have $E[S_t] = 0$, $E[S_t^2] = t\sigma^2$, and

$$\begin{aligned}\gamma_S(t+h, t) &= \text{Cov}(S_{t+h}, S_t) \\ &= \text{Cov}\left(S_t + \sum_{s=1}^h X_{t+s}, S_t\right) \\ &= \text{Cov}(S_t, S_t) = t\sigma^2.\end{aligned}$$

1. $\mu_t = 0$ is independent of t , but
2. $\gamma_S(t+h, t)$ is not.

So $\{S_t\}$ is not stationary.

An aside: covariances


$$\text{Cov}(X + Y, Z) = \text{Cov}(X, Z) + \text{Cov}(Y, Z),$$

$$\text{Cov}(aX, Y) = a \text{Cov}(X, Y),$$

Also if X and Y are independent (e.g., $X = c$), then

$$\text{Cov}(X, Y) = 0.$$

Random walk



Stationarity

Example: MA(1) process (**Moving Average**):


$$X_t = W_t + \theta W_{t-1}, \quad \{W_t\} \sim WN(0, \sigma^2).$$

We have $E[X_t] = 0$, and

$$\begin{aligned} \gamma_X(t+h, t) &= E(X_{t+h}X_t) \\ &= E[(W_{t+h} + \theta W_{t+h-1})(W_t + \theta W_{t-1})] \\ &= \begin{cases} \sigma^2(1 + \theta^2) & \text{if } h = 0, \\ \sigma^2\theta & \text{if } h = \pm 1, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Thus, $\{X_t\}$ is stationary.

ACF of the MA(1) process



Stationarity

Example: AR(1) process (**AutoRegressive**):

$$X_t = \phi X_{t-1} + W_t, \quad \{W_t\} \sim WN(0, \sigma^2).$$

Assume that X_t is stationary and $|\phi| < 1$. Then we have

$$\begin{aligned} \mathbb{E}[X_t] &= \phi \mathbb{E}X_{t-1} \\ &= 0 \quad (\text{from stationarity}) \end{aligned}$$


$$\begin{aligned} \mathbb{E}[X_t^2] &= \phi^2 \mathbb{E}[X_{t-1}^2] + \sigma^2 \\ &= \frac{\sigma^2}{1 - \phi^2} \quad (\text{from stationarity}), \end{aligned}$$

Stationarity

Example: AR(1) process, $X_t = \phi X_{t-1} + W_t$, $\{W_t\} \sim WN(0, \sigma^2)$.
Assume that X_t is stationary and $|\phi| < 1$. Then we have

$$\begin{aligned} E[X_t] &= 0, & E[X_t^2] &= \frac{\sigma^2}{1 - \phi^2} \\ \gamma_X(h) &= \text{Cov}(\phi X_{t+h-1} + W_{t+h}, X_t) \\ &= \phi \text{Cov}(X_{t+h-1}, X_t) \\ &= \phi \gamma_X(h-1) \\ &= \phi^{|h|} \gamma_X(0) && \text{(check for } h > 0 \text{ and } h < 0) \\ &= \frac{\phi^{|h|} \sigma^2}{1 - \phi^2}. \end{aligned}$$

ACF of the AR(1) process



Linear Processes

An important class of stationary time series:

$$X_t = \mu + \sum_{j=-\infty}^{\infty} \psi_j W_{t-j}$$

where $\{W_t\} \sim WN(0, \sigma_w^2)$

and μ, ψ_j are parameters satisfying

$$\sum_{j=-\infty}^{\infty} |\psi_j| < \infty.$$

Linear Processes

$$X_t = \mu + \sum_{j=-\infty}^{\infty} \psi_j W_{t-j}$$

We have

$$\mu_X = \mu$$

$$\gamma_X(h) = \sigma_w^2 \sum_{j=-\infty}^{\infty} \psi_j \psi_{h+j}. \quad (\text{why?})$$

Examples of Linear Processes: White noise

$$X_t = \mu + \sum_{j=-\infty}^{\infty} \psi_j W_{t-j}$$

Choose $\mu,$

$$\psi_j = \begin{cases} 1 & \text{if } j = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Then $\{X_t\} \sim WN(\mu, \sigma_W^2).$

(why?)

Examples of Linear Processes: MA(1)

$$X_t = \mu + \sum_{j=-\infty}^{\infty} \psi_j W_{t-j}$$

Choose $\mu = 0$

$$\psi_j = \begin{cases} 1 & \text{if } j = 0, \\ \theta & \text{if } j = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Then $X_t = W_t + \theta W_{t-1}$.

(why?)

Examples of Linear Processes: AR(1)

$$X_t = \mu + \sum_{j=-\infty}^{\infty} \psi_j W_{t-j}$$

Choose $\mu = 0$

$$\psi_j = \begin{cases} \phi^j & \text{if } j \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

Then for $|\phi| < 1$, we have $X_t = \phi X_{t-1} + W_t$.

(why?)

Introduction to Time Series Analysis. Lecture 2.

1. Stationarity
2. Autocovariance, autocorrelation
3. MA, AR, linear processes