

# **Introduction to Time Series Analysis. Lecture 3.**

**Peter Bartlett**

Last lecture:

1. Stationarity
2. Autocovariance, autocorrelation
3. MA, AR, linear processes

# **Introduction to Time Series Analysis. Lecture 3.**

**Peter Bartlett**

1. Review: Autocovariance, linear processes
2. Sample autocorrelation function
3. ACF and prediction
4. Properties of the ACF

## Mean, Autocovariance, Stationarity

A time series  $\{X_t\}$  has **mean function**  $\mu_t = E[X_t]$  and **autocovariance function**

$$\begin{aligned}\gamma_X(t+h, t) &= \text{Cov}(X_{t+h}, X_t) \\ &= E[(X_{t+h} - \mu_{t+h})(X_t - \mu_t)].\end{aligned}$$

It is **stationary** if both are independent of  $t$ .

Then we write  $\gamma_X(h) = \gamma_X(h, 0)$ .

The **autocorrelation function (ACF)** is

$$\rho_X(h) = \frac{\gamma_X(h)}{\gamma_X(0)} = \text{Corr}(X_{t+h}, X_t).$$

## Linear Processes

An important class of stationary time series:

$$X_t = \mu + \sum_{j=-\infty}^{\infty} \psi_j W_{t-j}$$

where  $\{W_t\} \sim WN(0, \sigma_w^2)$

and  $\mu, \psi_j$  are parameters satisfying

$$\sum_{j=-\infty}^{\infty} |\psi_j| < \infty.$$

## Linear Processes

$$X_t = \mu + \sum_{j=-\infty}^{\infty} \psi_j W_{t-j}$$

Examples:

- White noise:  $\psi_0 = 1$ .
- MA(1):  $\psi_0 = 1, \psi_1 = \theta$ .
- AR(1):  $\psi_0 = 1, \psi_1 = \phi, \psi_2 = \phi^2, \dots$

## Estimating the ACF: Sample ACF

Recall:

Suppose that  $\{X_t\}$  is a stationary time series.  
Its **mean** is

$$\mu = E[X_t].$$

Its **autocovariance function** is

$$\begin{aligned}\gamma(h) &= \text{Cov}(X_{t+h}, X_t) \\ &= E[(X_{t+h} - \mu)(X_t - \mu)].\end{aligned}$$

Its **autocorrelation function** is

$$\rho(h) = \frac{\gamma(h)}{\gamma(0)}.$$

## Estimating the ACF: Sample ACF

For observations  $x_1, \dots, x_n$  of a time series,

the **sample mean** is 
$$\bar{x} = \frac{1}{n} \sum_{t=1}^n x_t.$$

The **sample autocovariance function** is

$$\hat{\gamma}(h) = \frac{1}{n} \sum_{t=1}^{n-|h|} (x_{t+|h|} - \bar{x})(x_t - \bar{x}), \quad \text{for } -n < h < n.$$

The **sample autocorrelation function** is

$$\hat{\rho}(h) = \frac{\hat{\gamma}(h)}{\hat{\gamma}(0)}.$$

## Estimating the ACF: Sample ACF

Sample autocovariance function:


$$\hat{\gamma}(h) = \frac{1}{n} \sum_{t=1}^{n-|h|} (x_{t+|h|} - \bar{x})(x_t - \bar{x}).$$

$\approx$  the sample covariance of  $(x_1, x_{h+1}), \dots, (x_{n-h}, x_n)$ , except that

- we normalize by  $n$  instead of  $n - h$ , and
- we subtract the full sample mean.



## Sample ACF for white Gaussian (hence i.i.d.) noise




Red lines=c.i.

## Sample ACF


We can recognize the sample autocorrelation functions of many non-white (even non-stationary) time series.

<b>Time series:</b>	<b>Sample ACF:</b>
White	zero
Trend	Slow decay
Periodic	Periodic
MA( $q$ )	Zero for $ h  > q$
AR( $p$ )	Decays to zero exponentially

## Sample ACF: Trend



## Sample ACF: Trend



(why?)

## Sample ACF

**Time series:**

White

Trend

Periodic

MA( $q$ )

AR( $p$ )

**Sample ACF:**

zero


Slow decay

Periodic


Zero for  $|h| > q$

Decays to zero exponentially


## Sample ACF: Periodic



## Sample ACF: Periodic



## Sample ACF: Periodic







## Sample ACF

Time series:	Sample ACF:
White	zero
Trend	Slow decay
Periodic	Periodic
MA( $q$ )	Zero for $ h  > q$
AR( $p$ )	Decays to zero exponentially

## ACF: MA(1)




## Sample ACF: MA(1)




## Sample ACF

Time series:	Sample ACF:
White	zero
Trend	Slow decay
Periodic	Periodic
MA( $q$ )	Zero for $ h  > q$
AR( $p$ )	Decays to zero exponentially

## ACF: AR(1)





## Sample ACF: AR(1)



## **Introduction to Time Series Analysis. Lecture 3.**


1. Sample autocorrelation function
2. ACF and prediction
3. Properties of the ACF

## ACF and prediction





## ACF of a MA(1) process



## ACF and least squares prediction

Best least squares estimate of  $Y$  is  $EY$ :

$$\min_c E(Y - c)^2 = E(Y - EY)^2.$$

Best least squares estimate of  $Y$  given  $X$  is  $E[Y|X]$ :

$$\begin{aligned}\min_f E(Y - f(X))^2 &= \min_f E [E[(Y - f(X))^2|X]] \\ &= E [E[(Y - E[Y|X])^2|X]] \\ &= \text{var}[Y|X].\end{aligned}$$

Similarly, the best least squares estimate of  $X_{n+h}$  given  $X_n$  is  $f(X_n) = E[X_{n+h}|X_n]$ .

## ACF and least squares prediction

Suppose that  $X = (X_1, \dots, X_{n+h})$  is jointly Gaussian:

$$f_X(x) = \frac{1}{(2\pi)^{n/2} |\Sigma|^{1/2}} \exp \left( -\frac{1}{2} (x - \mu)' \Sigma^{-1} (x - \mu) \right).$$

Then the joint distribution of  $(X_n, X_{n+h})$  is

$$N \left( \begin{pmatrix} \mu_n \\ \mu_{n+h} \end{pmatrix}, \begin{pmatrix} \sigma_n^2 & \rho \sigma_n \sigma_{n+h} \\ \rho \sigma_n \sigma_{n+h} & \sigma_{n+h}^2 \end{pmatrix} \right),$$

and the conditional distribution of  $X_{n+h}$  given  $X_n$  is

$$N \left( \mu_{n+h} + \rho \frac{\sigma_{n+h}}{\sigma_n} (x_n - \mu_n), \sigma_{n+h}^2 (1 - \rho^2) \right).$$

## ACF and least squares prediction

So for Gaussian and stationary  $\{X_t\}$ , the best estimate of  $X_{n+h}$  given  $X_n = x_n$  is

$$f(x_n) = \mu + \rho(h)(x_n - \mu),$$

and the mean squared error is

$$E(X_{n+h} - f(X_n))^2 = \sigma^2(1 - \rho(h)^2).$$

Notice:

- Prediction accuracy improves as  $|\rho(h)| \rightarrow 1$ .
- Predictor is linear:  $f(x) = \mu(1 - \rho(h)) + \rho(h)x$ .

## ACF and least squares linear prediction

Consider a **linear predictor** of  $X_{n+h}$  given  $X_n = x_n$ . Assume first that  $\{X_t\}$  is stationary with  $EX_n = 0$ , and predict  $X_{n+h}$  with  $f(x_n) = ax_n$ . The best linear predictor minimizes

$$\begin{aligned} E(X_{n+h} - aX_n)^2 &= E(X_{n+h}^2) - E(2aX_{n+h}X_n) + E(a^2X_n^2) \\ &= \sigma^2 - 2a\gamma(h) + a^2\sigma^2, \end{aligned}$$

and this is minimized when  $a = \rho(h)$ , that is,

$$f(x_n) = \rho(h)X_n.$$

For this optimal linear predictor, the mean squared error is

$$\begin{aligned} E(X_{n+h} - f(X_n))^2 &= \sigma^2 - 2\rho(h)\gamma(h) + \rho(h)^2\sigma^2 \\ &= \sigma^2(1 - \rho(h)^2). \end{aligned}$$

## ACF and least squares linear prediction

Consider the following **linear predictor** of  $X_{n+h}$  given  $X_n = x_n$ , when  $\{X_n\}$  is stationary and  $EX_n = \mu$ :

$$f(x_n) = a(x_n - \mu) + b.$$

The linear predictor that minimizes

$$E(X_{n+h} - (a(X_n - \mu) + b))^2$$

has  $a = \rho(h)$ ,  $b = \mu$ , that is,

$$f(x_n) = \rho(h)(X_n - \mu) + \mu.$$

For this optimal linear predictor, the mean squared error is again

$$E(X_{n+h} - f(X_n))^2 = \sigma^2(1 - \rho(h)^2).$$

## Least squares prediction of $X_{n+h}$ given $X_n$

$$f(X_n) = \mu + \rho(h)(X_n - \mu).$$

$$E(f(X_n) - X_{n+h})^2 = \sigma^2(1 - \rho(h)^2).$$

- If  $\{X_t\}$  is stationary,  $f$  is the **optimal linear predictor**.
- If  $\{X_t\}$  is also Gaussian,  $f$  is the **optimal predictor**.
- Linear prediction is optimal for Gaussian time series.
- Over all stationary processes with that value of  $\rho(h)$  and  $\sigma^2$ , the optimal mean squared error is maximized by the Gaussian process.
- Linear prediction needs only second order statistics.
- Extends to longer histories,  $(X_n, X_{n-1}, \dots)$ .

## **Introduction to Time Series Analysis. Lecture 3.**

1. Sample autocorrelation function
2. ACF and prediction
3. Properties of the ACF



## Properties of the autocovariance function

For the autocovariance function  $\gamma$  of a stationary time series  $\{X_t\}$ ,

1.  $\gamma(0) \geq 0$ , (variance is non-negative)
2.  $|\gamma(h)| \leq \gamma(0)$ , (from Cauchy-Schwarz)
3.  $\gamma(h) = \gamma(-h)$ , (from stationarity)
4.  $\gamma$  is positive semidefinite.

Furthermore, any function  $\gamma : \mathbb{Z} \rightarrow \mathbb{R}$  that satisfies (3) and (4) is the autocovariance of some stationary time series.

## Properties of the autocovariance function

A function  $f : \mathbb{Z} \rightarrow \mathbb{R}$  is *positive semidefinite* if for all  $n$ , the matrix  $F_n$ , with entries  $(F_n)_{i,j} = f(i - j)$ , is positive semidefinite.

A matrix  $F_n \in \mathbb{R}^{n \times n}$  is positive semidefinite if, for all vectors  $a \in \mathbb{R}^n$ ,

$$a' F a \geq 0.$$

To see that  $\gamma$  is psd, consider the variance of  $(X_1, \dots, X_n)a$ .

## Properties of the autocovariance function

For the autocovariance function  $\gamma$  of a stationary time series  $\{X_t\}$ ,

1.  $\gamma(0) \geq 0$ ,
2.  $|\gamma(h)| \leq \gamma(0)$ ,
3.  $\gamma(h) = \gamma(-h)$ ,
4.  $\gamma$  is positive semidefinite.

Furthermore, any function  $\gamma : \mathbb{Z} \rightarrow \mathbb{R}$  that satisfies (3) and (4) is the autocovariance of some stationary time series (in particular, a Gaussian process).

e.g.: (1) and (2) follow from (4).

## **Introduction to Time Series Analysis. Lecture 3.**

1. Sample autocorrelation function
2. ACF and prediction
3. Properties of the ACF