

Optimal mean width and metric entropy estimates for convex bodies

(joint work w/ Grigoris Paouris)

Reese Pathak

UC Berkeley & Cornell University

September 29, 2026

The “size” of convex bodies

convex bodies: $K \subset \mathbb{R}^n$ is a *convex body* if it is compact, convex, and has nonempty interior

MAIN QUESTION OF THIS TALK

how “large” may a convex body $K \subset \mathbb{R}^n$ be, when its volume is fixed?

two notions of size:

1. *covering number* — how many ellipsoids of similar volume are required to cover K ?
2. *volume of projections* — what is the average size of a k -dimensional projection of K ?

(we will see that these two notions are, of course, very closely related)

Outline


1. Covering a convex body
2. Projections of a convex body
3. Proof ideas

1. Covering a convex body

Covering number and metric entropy

covering number: for compact sets $K, L \subset \mathbf{R}^n$ with L having nonempty interior,

$$N(K, L) = \min \left\{ m \geq 1 : \exists x_1, \dots, x_m \in \mathbf{R}^n \text{ such that } K \subset \bigcup_{i=1}^m x_i + L \right\}$$



metric entropy: logarithm of covering number, $\log N(K, L)$

Background on metric entropy and covering numbers

covering numbers:

- typically attributed to [Kolmogorov, '56; Kolmogorov-Tihomirov, '59], who wanted to understand how to succinctly approximate a function to a given level of accuracy
- appeared in metric space setting in [Pontrjagin-Schnirelmann, '32; von Neumann, '42]

entropy numbers and compactness of operators:

- given linear operator $u: X \rightarrow Y$ between Banach spaces, involves the study of

$$r \mapsto N(u(B_X), rB_Y), \quad r > 0,$$

where B_X, B_Y are unit balls of spaces X, Y

- leads to the *entropy numbers*, defined as

$$e_k(u) = \inf\{r > 0 : N(u(B_X), rB_Y) \leq 2^{k-1}\}, \quad k \geq 1$$

- quantifies compactness: note that u is compact iff $e_k \downarrow 0$ as $k \rightarrow \infty$
- typically attributed to [Pietsch, '72] though also appeared in [Mitjagin, '61; Mitjagin-Pelczynski, '68; Triebel, '70]

Background on metric entropy and covering numbers

existence of M -ellipsoids:

- [Milman, '86] established for convex bodies $K \subset \mathbf{R}^n$ which are centrally symmetric (i.e., $K = -K$) that there exists an ellipsoid $\mathcal{E}$ such that:

$$|\mathcal{E}| = |K| \quad \text{and} \quad N(K, \mathcal{E}) \leq \exp(Cn).$$

- [Milman-Pajor, 2000] obtained the same result in the non-symmetric case, up to translations

regular M -ellipsoids:

- [Pisier, '89a; Pisier, '89b] showed the existence of *regular* M -ellipsoids: for a centrally symmetric convex body $K \subset \mathbf{R}^n$ and each $\alpha \in (0, 2)$, there exists an ellipsoid $\mathcal{E}_\alpha$ such that

$$\max\{N(K, r\mathcal{E}_\alpha), N(\mathcal{E}_\alpha, rK), N(K^\circ, r\mathcal{E}_\alpha^\circ), N(\mathcal{E}_\alpha^\circ, rK^\circ)\} \leq \exp(C_\alpha n/r^\alpha), \quad \text{for } r \geq C_\alpha^{1/\alpha}$$

- weaker generalizations for non-symmetric bodies appeared in [Vritsiou, 2024]

duality of entropy:

- Pietsch's duality of entropy conjecture was partially resolved [Artstein-Milman-Szarek, 2004]:

$$N(\mathcal{E}^\circ, \tfrac{1}{c}K^\circ)^{1/C} \leq N(K, \mathcal{E}) \leq N(\mathcal{E}^\circ, cK^\circ)^C$$

for any centrally symmetric convex body $K \subset \mathbf{R}^n$, any ellipsoid $\mathcal{E}$

Covering numbers of convex bodies

THEOREM [PAOURIS-P., 2026]

for any convex body $K_0 \subset \mathbf{R}^n$, there is a translate K and ellipsoid $\mathcal{E}$ of the same volume such that

$$\begin{aligned} \max\{N(K, r\mathcal{E}), N(K^\circ, r\mathcal{E}^\circ), N(\mathcal{E}, rK), N(\mathcal{E}^\circ, rK^\circ)\} \\ \leq \inf_{|\tilde{\mathcal{E}}|=|B_1^n|} \max\{N(B_1^n, cr\tilde{\mathcal{E}}), N(B_\infty^n, cr\tilde{\mathcal{E}}^\circ), N(\tilde{\mathcal{E}}, crB_1^n), N(\tilde{\mathcal{E}}^\circ, crB_\infty^n)\}^C, \end{aligned}$$

holds for every $r > 0$

some remarks:

- infimum over ellipsoids $\tilde{\mathcal{E}}$ is attained (up to constants) with suitably scaled ball B_2^n
- the ellipsoid $\mathcal{E}$ does *not* depend on r
- result does *not* require symmetry of body K
- may replace crosspolytope B_1^n with any regular n -simplex S_n (after translation)

Equivalent formulation via entropy numbers

entropy numbers:

- *entropy numbers of a convex body* $K \subset \mathbf{R}^n$ are

$$e_k(K) = \inf\{r > 0 : N(K, rB_2^n) \leq 2^{k-1}\}, \quad e_k^*(K) = \inf\{r > 0 : N(B_2^n, rK^\circ) \leq 2^{k-1}\}$$

- for centrally symmetric K , these correspond to $e_k(\text{id})$, $e_k(\text{id}^*)$ for $\text{id}: (\mathbf{R}^n, \|\cdot\|_K) \rightarrow (\mathbf{R}^n, \|\cdot\|_2)$
- we set $E_k(K) = \max\{e_k(K), e_k^*(K), e_k(K^\circ), e_k^*(K^\circ)\}$

equivalent re-formulation of previous theorem:

THEOREM [PAOURIS-P., 2026]

for any convex body $K \subset \mathbf{R}^n$, there is an affine map ψ such that $|\psi(K)| = |B_2^n|$, and

$$E_k(\psi(K)) \lesssim \inf_{|\tilde{\psi}(B_1^n)| = |B_2^n|} E_k(\tilde{\psi}(B_1^n)), \quad \text{for all } k \geq 1$$

Comparison with the regular positions of Pisier

Pisier's extension of V. Milman's M -ellipsoid theorem [Pisier, '89a] can be formulated as follows

THEOREM [PISIER, '89A]

for any centrally symmetric convex body $K \subset \mathbf{R}^n$, any $\beta > 1/2$,
there exists $T_\beta \in \text{GL}(n)$ such that $|T_\beta K| = |B_2^n|$ and:

$$E_k(T_\beta K) \lesssim \max \left\{ 1, \frac{1}{\sqrt{2\beta - 1}} \right\} \left(\frac{n}{k} \right)^\beta, \quad \text{for } k \in \{1, 2, \dots, n\}$$

some remarks:

- fixing $k \leq n$ and then optimizing in β , Pisier's theorem implies

$$\inf_{|S_k K| = |B_2^n|} E_k(S_k K) \lesssim \sqrt{\frac{n}{k} \log \left(\frac{en}{k} \right)}$$

- in fact, essentially by [Schütt, '84], our theorem implies existence of $T \in \text{GL}(n)$

$$|TK| = |B_2^n|, \quad E_k(TK) \lesssim \min \left\{ \sqrt{n}, \sqrt{\frac{n}{k} \log \left(\frac{en}{k} \right)} \right\} \quad \text{simultaneously for } k \in \{1, 2, \dots, n\}$$

- our theorem also implies the above in non-symmetric case, after a translation

2. Projections of a convex body

Projections of a convex body: mean width and quermassintegrals

QUERMASSEINTEGRALS OF A CONVEX BODY

$$W_{[k]}(K) = \mathbf{E}_{F \sim \nu_{n,k}} \left[\frac{|P_F K|}{|B_2^k|} \right]^{1/k}, \quad \text{for } k \in \{1, 2, \dots, n\}$$

- $\nu_{n,k}$ denotes uniform (Haar) probability measure on all k -dimensional subspaces of $\mathbf{R}^n$
- P_F denotes orthogonal projection onto k -dimensional subspace $F \subset \mathbf{R}^n$

important special cases:

- for $k = 1$, this is the *spherical mean width*

$$W_{[1]}(K) = \mathbf{E}_{\theta \sim \sigma_{n-1}} \left[\sup_{x \in K} \langle x, \theta \rangle \right] \equiv M^*(K)$$

- for $k = n - 1$, this is the (scaled) *surface area*,

$$W_{[n-1]}(K) = \left(\frac{|\partial K|}{|\partial B_2^n|} \right)^{1/(n-1)}$$

- for $k = n$, this is the *volume radius*,

$$W_{[n]}(K) = \left(\frac{|K|}{|B_2^n|} \right)^{1/n} \equiv \text{vr}(K)$$

Background on mean width and quermassintegrals

- *Steiner formula*: for a convex body $K \subset \mathbf{R}^n$ and any $t \geq 0$,

$$|K + tB_2^n| = \sum_{k=0}^n |B_2^k| V_{n-k}(K) t^k$$

- coefficients $\{V_k(K)\}_{k=0}^n$ are referred to as *intrinsic volumes* and satisfy

$$V_k(K) = c_{n,k} \cdot W_{[k]}(K)^k \quad \text{where} \quad c_{n,k} = \binom{n}{k} \frac{|B_2^n|}{|B_2^{n-k}|}$$

- our normalization and ordering of the quermassintegrals are not entirely standard
- by Alexandrov inequalities, have

$$M^*(K) \equiv W_{[1]}(K) \geq W_{[2]}(K) \geq \cdots \geq W_{[n]}(K) \equiv \text{vr}(K)$$

- the inequality $M^*(K) \geq \text{vr}(K)$ is referred to as *Urysohn's inequality*
- extremizing $W_{[k]}(K)$ over bodies of fixed volume $\longleftrightarrow$ “generalized” isoperimetric problems

Background on mean width and quermassintegrals

- for $k = n - 1$, the classical isoperimetric theorem implies that

$$W_{[n-1]}(K) \geq \text{vr}(K),$$

with equality iff K is the ball $\text{vr}(K) B_2^n$ (up to translation)

- in the other direction, a theorem of [Ball, '91] shows

$$\inf_{T \in \text{SL}(n)} W_{[n-1]}(TK) \leq W_{[n-1]}(S_n), \quad \text{where } |K| = |S_n|,$$

and S_n is a regular n -simplex; equality holds iff K is a simplex [Barthe, '98]

- maximal order of the minimal k th quermassintegral, i.e.,

$$A_{n,k} = \sup_K \inf_{T \in \text{SL}(n)} \frac{W_{[k]}(TK)}{\text{vr}(K)}$$

was (previously) unknown for $k \in \{1, 2, \dots, cn\}$

- $k \gtrsim n$: can show $A_{n,k} \simeq 1$, via the work of [Milman, '86; Milman-Pajor, 2000]
- our methods, unfortunately, are unlikely to yield exact constants, equality cases

Volumes of typical k -dimensional projections of convex bodies

THEOREM [PAOURIS-P., 2026]

for any convex body $K_0 \subset \mathbf{R}^n$, there is a translate K and $T \in \mathrm{SL}(n)$ such that

$$\max \left\{ \frac{W_{[k]}(TK)}{\mathrm{vr}(K)}, \frac{W_{[k]}((TK)^\circ)}{\mathrm{vr}(K^\circ)} \right\} \lesssim \inf_{\tilde{T} \in \mathrm{SL}(n)} \max \left\{ \frac{W_{[k]}(\tilde{T}B_1^n)}{\mathrm{vr}(B_1^n)}, \frac{W_{[k]}((\tilde{T}B_1^n)^\circ)}{\mathrm{vr}(B_\infty^n)} \right\},$$

simultaneously, for every $k \in \{1, 2, \dots, n\}$

some remarks:

- essentially by a computation in [Paouris-Pivovarov, 2013], it holds that

$$\inf_{\tilde{T} \in \mathrm{SL}(n)} \max \left\{ \frac{W_{[k]}(\tilde{T}B_1^n)}{\mathrm{vr}(B_1^n)}, \frac{W_{[k]}((\tilde{T}B_1^n)^\circ)}{\mathrm{vr}(B_\infty^n)} \right\} \simeq \sqrt{\log \left(\frac{en}{k} \right)}$$

- important special case is *minimal mean width*: for any convex body $K \subset \mathbf{R}^n$,

$$1 \leq \inf_{T \in \mathrm{SL}(n)} \frac{M^*(TK)}{\mathrm{vr}(K)} \lesssim \sqrt{\log(en)},$$

- in all the results above, may replace B_1^n by a regular n -simplex, up to translation

History of minimal mean width

- the question about the order of the minimal mean width of convex bodies seems to have first appeared in [Giannopoulos-Milman-Rudelson, 2000]
- it is mentioned in the books [Aubrun-Szarek, 2017; Artstein-Avidan-Giannopoulos-Milman, 2015]
- following the introduction of the K -convexity $\mathbf{K}(K)$ constant, [Figiel-Tomczak-Jaegermann, '79] showed that in the centrally symmetric case,

$$\inf_{T \in \text{GL}(n)} M^*(TK) M(TK) \lesssim \mathbf{K}(K) \lesssim \log(1 + d_{\text{BM}}(K, B_2^n)) \lesssim \log(en)$$

- in the centrally symmetric case, as $M(TK) \gtrsim \text{vr}(TK)^{-1}$, it implies

$$\inf_{T \in \text{GL}(n)} \frac{M^*(TK)}{\text{vr}(K)} \lesssim \log(en),$$

for the non-symmetric case, also follows via Rogers-Shephard inequality

- [Bizeul, 2026] established maximal order of $M^*(K)/\text{vr}(K)$ among bodies K in isotropic position; also implies order of minimal mean width for general convex bodies

3. Proof ideas

Convex comparison of continuous martingales

the following result seems to date (at least) back to [Borkar, '87]

PROPOSITION

for any $\mathbf{R}^n$ -valued, continuous L^2 martingale $\{M_t\}_{0 \leq t \leq T}$ with $M_0 = 0$,
if $d[M]_t \geq \rho I_n dt$ for some $\rho > 0$, then, for every convex $F: \mathbf{R}^n \rightarrow \mathbf{R}$,

$$\mathbf{E} F(M_T) \geq \mathbf{E}_{G \sim \gamma_n} F\left(\sqrt{\rho T} G\right)$$

some remarks:

- we apply with $F = h_K$, for some convex body K containing the origin in its interior; in that case,

$$\sqrt{n} M^*(K) \lesssim \frac{1}{\sqrt{\rho T}} \cdot \mathbf{E} h_K(M_T) = \frac{1}{\sqrt{\rho T}} \cdot \mathbf{E} \|M_T\|_{K^\circ}$$

- we will arrange that the body and martingale $\{M_t\}_{t \geq 0}$ satisfy

$$|K^\circ|^{1/n} \simeq 1, \quad \mathbf{E} \|M_T\|_{K^\circ} \lesssim 1 \quad \text{for} \quad T \simeq \frac{1}{\log(en)}$$

and that the proposition holds with $\rho \gtrsim 1$

The stochastic localization process

our martingales are constructed via stochastic localization process [Eldan, 2012; Eldan, 2013; Lee-Vempala, 2024]

CONSTRUCTION OF STOCHASTIC LOCALIZATION PROCESS

given a centered probability measure μ on $\mathbf{R}^n$, study tilt measures

$$d\mu_{t,\eta} \propto e^{\langle x, \eta \rangle - t\|x\|_2^2/2} d\mu, \quad a_t(\eta) = \mathbf{E}_{X \sim \mu_{t,\eta}}[X], \quad A_t(\eta) = \text{Cov}_{X \sim \mu_{t,\eta}}(X)$$

construct the *random tilt process*,

$$d\theta_t = a_t(\theta_t)dt + dB_t, \quad \theta_0 = 0$$

leads to random measure, mean, covariance process:

$$\mu_t = \mu_{t,\theta_t}, \quad a_t = a_t(\theta_t), \quad A_t = A_t(\theta_t)$$

Interpretation of the stochastic localization process

filtering interpretation: (observed by [El Alaoui-Montanari, 2022; Klartag-Putterman, 2023])

- it holds that

$$\text{Law}\left(\{tX + B_t\}_{t \geq 0}\right) = \text{Law}\left(\{\theta_t\}_{t \geq 0}\right),$$

where $X \sim \mu$ independently of $\{B_t\}_{t \geq 0}$

- also, $\mu_t = \text{Law}(X \mid tX + B_t) = \text{Law}(X \mid X + t^{-1}B_t)$ for $t > 0$
- pathwise: essentially “interpolate” between μ (at $t \rightarrow 0^+$) and δ_X (as $t \rightarrow \infty$)

two key processes: for μ supported on (sufficiently smooth) convex body L ,

- *mean process:*

$$a_t = \int_L x p_t(x) \, dx = \int_0^t A_s \, dB_s, \quad d[a]_t = A_t^2 \, dt$$

- *normal process:*

$$a_t^\circ = \int_{\partial L} n_L(x) p_t(x) \, d\mathcal{H}^{n-1}(x) = \int_L \nabla \log p_t(x) p_t(x) \, dx$$

Bobkov's maximal Gaussian measure position

[Bobkov, 2011] defined a c.s. convex body $K \subset \mathbf{R}^n$ to be in *maximal Gaussian measure position* if

$$\gamma_n(K) \geq \gamma_n(TK) \quad \text{for all } T \in \text{SL}(n)$$

GENERALIZATION TO NON-SYMMETRIC CASE ('B-POSITION')

a (possibly non-symmetric) convex body $K \subset \mathbf{R}^n$ is in *maximal Gaussian measure position* if

$$\gamma_n(K) \geq \gamma_n(TK + x) \quad \text{for all } (T, x) \in \text{SL}(n) \times \mathbf{R}^n$$

properties:

- equivalent characterization via isotropy: if γ_K denotes γ_n conditioned to K

$$\mathbf{E}_{X \sim \gamma_K} [X] = 0 \quad \text{and} \quad \mathbf{E}_{X \sim \gamma_K} [X \otimes X] = \alpha_K^2 I_n$$

- if $|K|^{1/n} \simeq 1$, then B-position is Milman's position
- for a convex body $K \subset \mathbf{R}^n$ there always exists a volume-preserving affine image of K which is in B-position; unique up to orthogonal transformations

A non-symmetric variant of the strong Gaussian (B)-theorem

the properties on the last slide can be established from the following result

PROPOSITION [PAOURIS-P., 2026]

for any real, symmetric $A \in \mathbf{R}^{n \times n}$ and any convex body $K \subset \mathbf{R}^n$, the map

$$t \mapsto \sup_{x \in \mathbf{R}^n} \gamma_n(e^{tA}K + x)$$

is log-concave on $\mathbf{R}$

some remarks:

- the result above is false without optimizing over translates [Nayar-Tkocz, 2013]
- proof of the result:
 - compute second derivative of the logarithm, realize it is nonpositive iff

$$\mathrm{Var}_{\gamma_K}(f) \leq \frac{1}{2} \left(\mathbf{E}_{\gamma_K} \|\nabla f\|_2^2 + \|\mathbf{E}_{\gamma_K} \nabla f\|_2^2 \right), \quad \text{for certain } f;$$

- follows by adapting proof of [Cordero-Erausquin-Fradelizi-Maurey, 2004; Barthe-Cordero-Erausquin, 2013]

Comparison theorem and width of K

THEOREM

for any convex body K with K° in B-position with $|K^\circ| = 1$,

$$\mathbf{E}_{\gamma_n} F(G) \lesssim \inf_{u \in (0, n]} \left(\sqrt{\log(en/u)} \mathbf{E}_{X \sim \gamma_{K^\circ}} F(X) + \text{Lip}(F) \sqrt{u} \right),$$

for any positively 1-homogeneous, convex $F: \mathbf{R}^n \rightarrow \mathbf{R}_+$

comments:

- we replace $da_t = A_t dB_t$ by $d\tilde{a}_t = \tilde{A}_t dB_t$ with $\tilde{A}_t$ explicitly truncated from below
- **the first term**: arises from the convex comparison for continuous martingales
- **the error term** $\text{Lip}(F) \sqrt{u}$: by time $T \simeq (\log(en/u))^{-1}$, have $\approx u$ truncated eigenvalues
- implies that

$$\sup_{x \in K} M^* \left(K \cap (x + r \text{vr}(K) B_2^n) \right) \lesssim \text{vr}(K) \sqrt{\log(er^2)} \quad (\forall r \geq 1)$$

Comparison theorem and width of K°

THEOREM

for any convex body K with K° in B-position with $|K^\circ| = 1$,

$$\mathbf{E}_{\gamma_n} F(G) \lesssim \inf_{u \in (0, n]} \left(n \sqrt{\log(en/u)} \mathbf{E}_{\nu_{K^\circ}} F(\nabla h_K(X)) + \text{Lip}(F) \sqrt{u} \right),$$

for any positively 1-homogeneous, convex $F: \mathbf{R}^n \rightarrow \mathbf{R}_+$.

(Here ν_{K° is the probability measure on ∂K° with $d\nu_{K^\circ}(x) \propto \langle x, n_{K^\circ}(x) \rangle e^{-\|x\|_2^2/2} d\mathcal{H}^{n-1}(x)$)

comments:

- the normal process satisfies $da_t^\circ = [I_n - (1+t)A_t]dB_t$; we now truncate eigenvalues of $(1+t)A_t$ from above
- **the first term**: arises from convex comparison, factor n comes from isotropy and volume normalization
- **the error term** $\text{Lip}(F)\sqrt{u}$: up to $T \simeq (\log(en/u))^{-1}$, have $\approx u$ truncated eigenvalues
- implies that

$$\sup_{x \in K^\circ} M^* \left(K^\circ \cap (x + r \text{vr}(K^\circ) B_2^n) \right) \lesssim \text{vr}(K^\circ) \sqrt{\log(er^2)} \quad (\forall r \geq 1)$$

Reduction to the truncation bodies

truncation bodies: for a convex body $L \subset \mathbf{R}^n$, we refer to

$$L \cap (x + rB_2^n), \quad \text{for } x \in L, r > 0$$

as the *truncation bodies* of L

PROPOSITION [PAOURIS-P., 2026]

suppose that $L \subset \mathbf{R}^n$ is a convex body with the origin in its interior, and with

$$\text{diam}(L) \lesssim \sqrt{n} \text{vr}(L), \quad \text{and} \quad \sup_{x \in L} M^*(L \cap (x + r \text{vr}(L) B_2^n)) \lesssim \text{vr}(L) \sqrt{\log(er^2)} \quad (\forall r \geq 1)$$

then for every $t > 0$,

$$\max \left\{ N(L, t \text{vr}(L) B_2^n), N(B_2^n, t \text{vr}(L) L^\circ) \right\} \leq \inf_{T \in \text{SL}(n)} N(TB_1^n, c \frac{t}{\sqrt{n}} B_2^n)^C$$

moreover, for every $k \in \{1, 2, \dots, n\}$

$$W_{[k]}(L) \lesssim \sqrt{n} \text{vr}(L) \inf_{T \in \text{SL}(n)} W_{[k]}(TB_1^n)$$

Ideas behind reduction

bounding the metric entropy:

- by applying the Sudakov minoration to “local” packing generated by a $t/2$ -packing and t -cover,

$$\log \frac{N(L, \frac{t}{2} B_2^n)}{N(L, t B_2^n)} \lesssim \frac{n}{t^2} \sup_{x \in L} M^*(L \cap (x + t B_2^n))^2$$

- by invoking the dual Sudakov minoration, we obtain

$$\log \frac{N(B_2^n, \frac{t}{2} L^\circ)}{N(B_2^n, t L^\circ)} \lesssim \frac{n}{t^2} \sup_{x \in L} M^*(L \cap (x + t B_2^n))^2$$

bounding the quermassintegrals:

THEOREM [MOURTADA, 2025]

for any convex body $K \subset \mathbf{R}^n$,

$$\log \left(\sum_{j=0}^n V_j(K) \right) \simeq \inf_{r>0} \left\{ \sqrt{n} \sup_{x \in K} M^*(K \cap (x + r B_2^n)) + \log N(K, r B_2^n) \right\}$$

- we apply this result to dilates λK ; optimize $\lambda > 0$ so that $V_k(\lambda_k K)$ dominates the sum

Summary

1. put K° in *B-position* such that $|K^\circ| = 1$
 - exists in non-symmetric case (by variant of strong (B) theorem)
 - equivalent to centered & scalar covariance for γ_{K°
 - implies that

$$\int_{K^\circ} x \, d\gamma_{K^\circ}(x) = 0, \quad \text{and} \quad \int_{\partial K^\circ} n_{K^\circ}(x) \phi(x) \, d\mathcal{H}^{n-1}(x) = 0$$

2. consider the continuous martingales

$$a_t = \mathbf{E}_{\mu_t} X, \quad \text{and} \quad a_t^\circ = \int_{\partial K^\circ} n_{K^\circ}(x) p_t(x) \, d\mathcal{H}^{n-1}(x)$$

3. by Itô computation,

$$da_t = A_t \, dB_t, \quad \text{and} \quad da_t^\circ = [I - (1+t)A_t] \, dB_t,$$

to apply martingale theorem requires controlling eigenvalues of covariance A_t , above and below

4. after controlling eigenvalues, apply *convex comparison theorem for continuous martingales*
5. take convex map to be *support function for truncated bodies* to get estimate for mean width
6. *reduce from mean width of truncated bodies* to metric entropy/quermassintegrals

References

References I

- Artstein, S., V. Milman, and S. J. Szarek (2004). “Duality of metric entropy”. In: *Ann. of Math. (2)* 159.3, pp. 1313–1328. ISSN: 0003-486X, 1939-8980. DOI: 10.4007/annals.2004.159.1313. URL: <https://doi.org/10.4007/annals.2004.159.1313>.
- Artstein-Avidan, Shiri, Apostolos Giannopoulos, and Vitali D. Milman (2015). *Asymptotic geometric analysis. Part I*. Vol. 202. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, pp. xx+451. ISBN: 978-1-4704-2193-9. DOI: 10.1090/surv/202. URL: <https://doi.org/10.1090/surv/202>.
- Aubrun, Guillaume and Stanisław J. Szarek (2017). *Alice and Bob meet Banach*. Vol. 223. Mathematical Surveys and Monographs. The interface of asymptotic geometric analysis and quantum information theory. American Mathematical Society, Providence, RI, pp. xxi+414. ISBN: 978-1-4704-3468-7. DOI: 10.1090/surv/223. URL: <https://doi.org/10.1090/surv/223>.
- Ball, Keith (1991). “Volume ratios and a reverse isoperimetric inequality”. In: *J. London Math. Soc. (2)* 44.2, pp. 351–359. ISSN: 0024-6107, 1469-7750. DOI: 10.1112/jlms/s2-44.2.351. URL: <https://doi.org/10.1112/jlms/s2-44.2.351>.
- Barthe, F. and D. Cordero-Erausquin (2013). “Invariances in variance estimates”. In: *Proc. Lond. Math. Soc. (3)* 106.1, pp. 33–64. ISSN: 0024-6115, 1460-244X. DOI: 10.1112/plms/pds011. URL: <https://doi.org/10.1112/plms/pds011>.

References II

- Barthe, Franck (1998). “On a reverse form of the Brascamp-Lieb inequality”. In: *Invent. Math.* 134.2, pp. 335–361. ISSN: 0020-9910,1432-1297. DOI: 10.1007/s002220050267. URL: <https://doi.org/10.1007/s002220050267>.
- Bizeul, Pierre (2026). *Optimal MM^* bounds for convex bodies*. arXiv: 2607.29458 [math.MG]. URL: <https://arxiv.org/abs/2607.29458>.
- Bobkov, Sergey G. (2011). “On Milman’s ellipsoids and M -position of convex bodies”. In: *Concentration, functional inequalities and isoperimetry*. Vol. 545. Contemp. Math. Amer. Math. Soc., Providence, RI, pp. 23–33. ISBN: 978-0-8218-4971-2. DOI: 10.1090/conm/545/10762. URL: <https://doi.org/10.1090/conm/545/10762>.
- Borkar, Vivek S. (1987). “A comparison principle for certain convex functionals of a diffusion process without drift”. In: *Stochastic Process. Appl.* 25.2, pp. 245–248. ISSN: 0304-4149,1879-209X. DOI: 10.1016/0304-4149(87)90201-8. URL: [https://doi.org/10.1016/0304-4149\(87\)90201-8](https://doi.org/10.1016/0304-4149(87)90201-8).
- Cordero-Erausquin, D., M. Fradelizi, and B. Maurey (2004). “The (B) conjecture for the Gaussian measure of dilates of symmetric convex sets and related problems”. In: *J. Funct. Anal.* 214.2, pp. 410–427. ISSN: 0022-1236,1096-0783. DOI: 10.1016/j.jfa.2003.12.001. URL: <https://doi.org/10.1016/j.jfa.2003.12.001>.
- El Alaoui, Ahmed and Andrea Montanari (2022). “An information-theoretic view of stochastic localization”. In: *IEEE Trans. Inform. Theory* 68.11, pp. 7423–7426. ISSN: 0018-9448,1557-9654. DOI: 10.1109/tit.2022.3180298. URL: <https://doi.org/10.1109/tit.2022.3180298>.

References III

- Eldan, Ronen (2012). “Distribution of Mass in Convex Bodies”. PhD thesis. Tel Aviv University.
- — (2013). “Thin shell implies spectral gap up to polylog via a stochastic localization scheme”. In: *Geom. Funct. Anal.* 23.2, pp. 532–569. ISSN: 1016-443X, 1420-8970. DOI: 10.1007/s00039-013-0214-y. URL: <https://doi.org/10.1007/s00039-013-0214-y>.
- Figiel, T. and Nicole Tomczak-Jaegermann (1979). “Projections onto Hilbertian subspaces of Banach spaces”. In: *Israel J. Math.* 33.2, pp. 155–171. ISSN: 0021-2172. DOI: 10.1007/BF02760556. URL: <https://doi.org/10.1007/BF02760556>.
- Giannopoulos, A. A., V. D. Milman, and M. Rudelson (2000). “Convex bodies with minimal mean width”. In: *Geometric aspects of functional analysis*. Vol. 1745. Lecture Notes in Math. Springer, Berlin, pp. 81–93. ISBN: 3-540-41070-8. DOI: 10.1007/BFb0107209. URL: <https://doi.org/10.1007/BFb0107209>.
- Klartag, Bo’az and Eli Putterman (2023). “Spectral monotonicity under Gaussian convolution”. In: *Ann. Fac. Sci. Toulouse Math. (6)* 32.5, pp. 939–967. ISSN: 0240-2963, 2258-7519. DOI: 10.5802/afst.1759. URL: <https://doi.org/10.5802/afst.1759>.
- Kolmogorov, A. N. (1956). “On certain asymptotic characteristics of completely bounded metric spaces”. In: *Dokl. Akad. Nauk SSSR (N.S.)* 108, pp. 385–388.
- Kolmogorov, A. N. and V. M. Tihomirov (1959). “ ε -entropy and ε -capacity of sets in function spaces”. In: *Uspehi Mat. Nauk* 14.2(86), pp. 3–86. ISSN: 0042-1316.

References IV

- Lee, Yin Tat and Santosh S. Vempala (2024). “Eldan’s stochastic localization and the KLS conjecture: isoperimetry, concentration and mixing”. In: *Ann. of Math. (2)* 199.3, pp. 1043–1092. ISSN: 0003-486X, 1939-8980. DOI: 10.4007/annals.2024.199.3.2. URL: <https://doi.org/10.4007/annals.2024.199.3.2>.
- Milman, V. D. and A. Pajor (2000). “Entropy and asymptotic geometry of non-symmetric convex bodies”. In: *Adv. Math.* 152.2, pp. 314–335. ISSN: 0001-8708, 1090-2082. DOI: 10.1006/aima.1999.1903. URL: <https://doi.org/10.1006/aima.1999.1903>.
- Milman, Vitali D. (1986). “Inégalité de Brunn-Minkowski inverse et applications à la théorie locale des espaces normés”. In: *C. R. Acad. Sci. Paris Sér. I Math.* 302.1, pp. 25–28. ISSN: 0249-6291.
- Mitjagin, B. S. (1961). “Approximate dimension and bases in nuclear spaces”. In: *Uspehi Mat. Nauk* 16.4(100), pp. 63–132. ISSN: 0042-1316.
- Mitjagin, B. S. and A. Pelczynski (1968). “Nuclear operators and approximative dimension”. In: *Proc. Internat. Congr. Math. (Moscow, 1966)*. Izdat. “Mir”, Moscow, pp. 366–372.
- Nayar, Piotr and Tomasz Tkocz (2013). “A note on a Brunn-Minkowski inequality for the Gaussian measure”. In: *Proc. Amer. Math. Soc.* 141.11, pp. 4027–4030. ISSN: 0002-9939, 1088-6826. DOI: 10.1090/S0002-9939-2013-11609-6. URL: <https://doi.org/10.1090/S0002-9939-2013-11609-6>.
- Paouris, Grigoris and Peter Pivovarov (2013). “Small-ball probabilities for the volume of random convex sets”. In: *Discrete Comput. Geom.* 49.3, pp. 601–646. ISSN: 0179-5376, 1432-0444. DOI: 10.1007/s00454-013-9492-2. URL: <https://doi.org/10.1007/s00454-013-9492-2>.

References V

- Pietsch, Albrecht (1972). *Theorie der Operatorenideale (Zusammenfassung)*. Wissenschaftliche Beiträge der Friedrich-Schiller-Universität Jena. Friedrich-Schiller-Universität, Jena, pp. x+260.
- Pisier, Gilles (1989a). “A new approach to several results of V. Milman”. In: *J. Reine Angew. Math.* 393, pp. 115–131. ISSN: 0075-4102,1435-5345. DOI: 10.1515/crll.1989.393.115. URL: <https://doi.org/10.1515/crll.1989.393.115>.
- — (1989b). *The volume of convex bodies and Banach space geometry*. Vol. 94. Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, pp. xvi+250. ISBN: 0-521-36465-5. DOI: 10.1017/CB09780511662454. URL: <https://doi.org/10.1017/CB09780511662454>.
- Pontrjagin, L. and L. Schnirelmann (1932). “Sur une propriété métrique de la dimension”. In: *Ann. of Math. (2)* 33.1, pp. 156–162. ISSN: 0003-486X,1939-8980. DOI: 10.2307/1968109. URL: <https://doi.org/10.2307/1968109>.
- Schütt, Carsten (1984). “Entropy numbers of diagonal operators between symmetric Banach spaces”. In: *J. Approx. Theory* 40.2, pp. 121–128. ISSN: 0021-9045,1096-0430. DOI: 10.1016/0021-9045(84)90021-2. URL: [https://doi.org/10.1016/0021-9045\(84\)90021-2](https://doi.org/10.1016/0021-9045(84)90021-2).
- Triebel, Hans (1970). “Interpolationseigenschaften von Entropie- und Durchmesseridealen kompakter Operatoren”. In: *Studia Math.* 34, pp. 89–107. ISSN: 0039-3223,1730-6337. DOI: 10.4064/sm-34-1-89-107. URL: <https://doi.org/10.4064/sm-34-1-89-107>.
- von Neumann, John (1942). “Approximative properties of matrices of high finite order”. In: *Portugal. Math.* 3, pp. 1–62. ISSN: 0032-5155,1662-2758.

References VI

- Vritsiou, Beatrice-Helen (2024). “Regular ellipsoids and a Blaschke-Santaló-type inequality for projections of non-symmetric convex bodies”. In: *J. Funct. Anal.* 286.11, Paper No. 110414, 40. ISSN: 0022-1236,1096-0783. DOI: 10.1016/j.jfa.2024.110414. URL: <https://doi.org/10.1016/j.jfa.2024.110414>.